

---

## A Grower Survey of Herbicide Use Patterns in Glyphosate-Resistant Cropping Systems

Author(s): Wade A. Givens, David R. Shaw, William G. Johnson, Stephen C. Weller, Bryan G. Young, Robert G. Wilson, Micheal D. K. Owen, and David Jordan

Source: Weed Technology, 23(1):156-161. 2009.

Published By: Weed Science Society of America

DOI: <http://dx.doi.org/10.1614/WT-08-039.1>

URL: <http://www.bioone.org/doi/full/10.1614/WT-08-039.1>

---

BioOne ([www.bioone.org](http://www.bioone.org)) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/page/terms\\_of\\_use](http://www.bioone.org/page/terms_of_use).

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

## A Grower Survey of Herbicide Use Patterns in Glyphosate-Resistant Cropping Systems

Wade A. Givens, David R. Shaw, William G. Johnson, Stephen C. Weller, Bryan G. Young, Robert G. Wilson, Micheal D. K. Owen, and David Jordan\*

A telephone survey was conducted with growers in Iowa, Illinois, Indiana, Nebraska, Mississippi, and North Carolina to discern the utilization of the glyphosate-resistant (GR) trait in crop rotations, weed pressure, tillage practices, herbicide use, and perception of GR weeds. This paper focuses on survey results regarding herbicide decisions made during the 2005 cropping season. Less than 20% of the respondents made fall herbicide applications. The most frequently used herbicides for fall applications were 2,4-D and glyphosate, and these herbicides were also the most frequently used for preplant burndown weed control in the spring. Atrazine and acetochlor were frequently used in rotations containing GR corn. As expected, crop rotations using a GR crop had a high percentage of respondents that made one to three POST applications of glyphosate per year. GR corn, GR cotton, and non-GR crops had the highest percentage of growers applying non-glyphosate herbicides during the 2005 growing season. A crop rotation containing GR soybean had the greatest negative impact on non-glyphosate use. Overall, glyphosate use has continued to increase, with concomitant decreases in utilization of other herbicides.

**Nomenclature:** 2,4-D; acetochlor; atrazine; glyphosate; corn, *Zea mays* L.; cotton, *Gossypium hirsutum* L.; soybean, *Glycine max* (L.) Merr.

**Key words:** Postemergence, preemergence, preplant burndown, soil residual herbicides, tank mixtures, weed management.

The introduction of 2,4-D in the mid-1940s ushered in a new era in which growers had a viable alternative to mechanical control of weeds (Burnside 1996). Over the following decades, there was an explosion of herbicide discovery that changed the way growers managed weeds. During this time of herbicide discovery, several nonselective herbicides were commercialized, including paraquat, glufosinate, and glyphosate.

Glyphosate was introduced to the market in the early 1970s for broad spectrum weed control, including perennial weeds. Glyphosate has become one of the world's leading agrochemicals (Woodburn 2000). During the 1970s and early 1980s, research explored means of breeding herbicide resistance into crops (Barrentine et al. 1982). However, it was not until the 1980s that the tools for developing genetically engineered, transgenic crops became available. Several companies saw the advantage of using these technologies to produce transgenic crops that would be resistant to herbicides. Extensive efforts were put forth to develop GR crops, eventually leading to the use of the CP4 gene from *Agrobacterium* sp. This gene codes for a glyphosate-insensitive 5-enol-pyruvylshikimate-3-phosphate synthase (EPSPS) in selected crops (Padgett et al. 1995).

The first commercially available GR crop was soybean, introduced in 1996. GR cotton followed in 1997, and GR corn was introduced in 1998. In 2007, 91% of soybean, 70%

of cotton, and 52% of the corn hectares were planted to GR cultivars in the United States (USDA-NASS 2007). Adoption of GR technologies has been rapid due to a wider spectrum of weeds controlled, less need for tank-mixing other herbicides, and reduced time and labor inputs (Ateh and Harvey 1999; Bradley et al. 2004; Corbett et al. 2004; Faircloth et al. 2001; Johnson et al. 2000; Reddy and Whiting 2000; Thomas et al. 2004a, b).

After nearly a decade of growing GR crops one would expect significant changes in herbicide use in terms of the frequency and amount of use for herbicide active ingredients. Several researchers have investigated herbicide use patterns following GR crop adoption by examining existing datasets such as the National Agricultural Statistics Service (NASS) chemical use databases and other industry compiled databases (Shaner 2000; Young 2006). An overall reduction in the amount of herbicides applied was noted since grower adoption of GR cropping systems, as was an increased reliance on glyphosate in their weed management programs.

One means of collecting data on actual usage and grower perceptions about weed management is through grower surveys. These surveys have been used in the past to document changes in management practices and grower perceptions about potential problems in a wide range of areas from irrigation practices to perceptions about insect pressure and pesticide use (Dillard 1993; Snyder 1996). Grower surveys have been especially important to weed science and have allowed scientists to gain insight into a number of grower perceptions and practices. Examples include herbicide and herbicide-resistant crop use and grower perceptions of issues such as herbicide resistance in weeds (Charles 1991; Gibson et al. 2005; Gibson et al. 2006; Johnson and Gibson 2006; Llewellyn et al. 2002). By using grower surveys, we have the opportunity to capture a cross-section of weed management

DOI: 10.1614/WT-08-039.1

\* Research Associate and Giles Distinguished Professor, Department of Plant and Soil Sciences, Mississippi State University, Mississippi State, MS 39762; Associate Professor and Professor, Purdue University, West Lafayette, IN 47907; Professor, Southern Illinois University, Carbondale, IL 62901; Professor, University of Nebraska, Scottsbluff, NE 69361; Professor, Iowa State University, Ames, IA 50011; and Professor, North Carolina State University, Raleigh, NC 27606. Corresponding author's E-mail: wgivens@gsi.msstate.edu

Table 1. Herbicide application summary for all reported crop rotations in Iowa, Illinois, Indiana, Mississippi, North Carolina, and Nebraska for the 2005 growing season. The responses are categorized by application timing.

Crop rotation	Fall application of herbicides	Burndown application prior to or at planting in the spring	Glyphosate applications				Non-glyphosate applications				Avg. no. yrs. an herbicide other than glyphosate was applied <sup>b</sup>
			1 application	2 applications	> 3 applications	Did not apply	Prior to planting	At planting	Post emerge	Did not apply	
% of growers making each application											
Continuous GR <sup>a</sup> corn (n = 84)	13	27	54	42	2	1	12	16	24	57	1.3
Continuous GR soybean (n = 307)	14	60	23	62	12	2	4	1	10	85	0.7
Continuous GR cotton (n = 97)	10	76	12	44	42	—	10	10	69	36	2.1
GR corn / GR soybean											
GR corn (n = 407)	7	36	63	32	2	3	18	20	13	56	1.2
GR soybean (n = 407)	9	38	48	47	3	2	9	2	6	84	0.9
GR cotton / GR Soybean											
GR cotton (n = 38)	16	76	18	47	29	—	3	5	58	47	1.6
GR soybean (n = 38)	8	63	26	53	13	3	5	—	16	79	1.1
GR soybean/on-GR crop											
GR soybean (n = 496)	8	42	52	43	4	1	11	2	8	81	1.3
Non-GR crop (n = 496)	6	27	—	—	—	—	22	46	53	23	1.9
GR corn / non-GR crop											
GR corn (n = 85)	14	25	61	31	2	6	17	25	20	45	1.8
Non-GR crop (n = 85)	4	20	—	—	—	—	12	31	53	33	1.5

<sup>a</sup> Abbreviations: GR, glyphosate-resistant.

<sup>b</sup> Average number of years non-glyphosate herbicide was applied (out of the last 3 yr).

practices and their potential problems spanning different states and crop rotations after using a GR crop.

The purpose of this paper is to determine and quantify the effect of GR crop use on growers' herbicide use patterns. The data for this paper are a subset from a dataset generated from a telephone survey that was conducted between November 9, 2005 and January 6, 2006, to capture many aspects of long-term GR crop use and the changes over time that have occurred because of their use.

## Materials and Methods

A survey was designed by the authors and conducted in six states. The telephone survey was conducted by contacting growers from Iowa, Illinois, Indiana, Mississippi, North Carolina, and Nebraska. A list of all growers from these states who had signed an agreement to use the glyphosate-resistant crop [Roundup Ready™] technology was obtained from the company, and survey respondents were randomly selected from this list. Respondents were initially asked whether they were actively involved in farming, if they were responsible for the management decisions in their farming operations, if they planted a minimum of 101 hectares of corn, soybean, or cotton in 2005, and if they planted one of the traits or trait combinations for a minimum of 3 yr. Producers were disqualified from the survey if they or anyone in their household worked for a farm chemical manufacturer, distributor, or retailer, or if they worked for a seed company other than as a farmer/dealer. The survey consisted of four sections dealing with different aspects of their farming practices. The sections dealt specifically with cropping history, weed pressure and tillage practices, herbicide use, and GR weeds. The respondents were asked to focus their answers on

one specific representative field for each cropping system. Complete details of the survey are reported in an introductory paper for this series by Shaw et al. (2009). This paper will focus mainly on the herbicide use data generated from the survey.

For this analysis, only the following crop rotations are discussed: continuous GR corn, continuous GR soybean, continuous GR cotton, GR corn / GR soybean rotation, GR cotton / GR soybean rotation, GR soybean / non-GR crop rotation, and GR corn / non-GR crop rotation. Grower responses on herbicide application timing and frequency within each cropping system are located in Table 1. Table 2 lists the most frequently used herbicide active ingredients for fall applications. Table 3 lists the herbicide active ingredients used for preplant burndown applications. Data presented in Table 4 are the applications of non-glyphosate herbicide active ingredients pooled across application timings and the percentage of growers in each crop rotation that did not apply an herbicide other than glyphosate during the cropping season in question.

## Results and Discussion

The data presented in Table 1 are a summary of responses to the questions relating to herbicide use. The data are categorized by crop rotation and herbicide system. The crop rotations examined included continuous GR corn, continuous GR cotton, continuous GR soybean, GR corn / GR soybean, GR cotton / GR soybean, GR soybean / non-GR crop, and GR corn / non-GR crop. The herbicide systems were broken out by fall application, preplant burndown application, glyphosate-in-crop applications, and non-glyphosate-in-crop applications. Glyphosate applications were further categorized

Table 2. Fall herbicides applied for each crop rotation. Data expressed as percentages of producers in each crop rotation who applied each herbicide.

Crop rotation	Atrazine	Simazine	Chlorimuron	Flumioxazin	Glyphosate	2,4-D	Dicamba	Paraquat
	% of growers making each application							
Continuous GR <sup>a</sup> corn ( <i>n</i> = 11)	—	—	—	—	36	9	—	—
Continuous GR soybean ( <i>n</i> = 43)	—	—	5	—	56	7	5	5
Continuous GR cotton ( <i>n</i> = 10)	—	—	—	—	30	—	—	—
GR corn / GR soybean								
GR corn ( <i>n</i> = 30)	10	17	3	—	16	13	—	—
GR soybean ( <i>n</i> = 37)	—	3	27	—	27	19	—	—
GR cotton / GR soybean								
GR cotton ( <i>n</i> = 0)	—	—	—	—	—	—	—	—
GR soybean ( <i>n</i> = 0)	—	—	—	—	—	—	—	—
GR soybean / non-GR crop								
GR soybean ( <i>n</i> = 41)	—	2	37	5	32	32	—	—
Non-GR crop ( <i>n</i> = 31)	10	26	—	—	10	16	—	—
GR corn / non-GR crop								
GR corn ( <i>n</i> = 12)	17	—	—	—	25	25	—	—
Non-GR crop ( <i>n</i> = 0)	—	—	—	—	—	—	—	—

<sup>a</sup> Abbreviations: GR, glyphosate-resistant.

by number of applications, and non-glyphosate applications were further categorized by timing of the applications. Data from each application timing category are discussed below.

**Fall Herbicide Use.** Between 4 and 16% of growers made fall applications of herbicides prior to planting the specified crop in 2005 (Table 1). Four to 6% of the respondents indicated they used a fall herbicide application prior to planting a non-GR crop. Conversely, at least 10% of the growers with crop rotations that included continuous GR corn, continuous GR soybean, and continuous GR cotton used a fall herbicide application. Thus, the use of fall herbicide application may be more common in continuous GR cropping systems. The cause of the increased need for fall herbicide applications in continuous GR cropping systems is beyond the scope of this survey. However, greater reliance on glyphosate and non-residual herbicides has been associated with greater problems with winter annual weeds. The most commonly used herbicides across all crop rotations were glyphosate and 2,4-D (Table 2). Atrazine, chlorimuron, and simazine were also frequently used herbicides, but their usage was very specific,

based on crop tolerances of each rotation. These herbicides are often applied in the fall to control weeds that would otherwise be difficult to manage in the spring and potentially compete with the crop (Wicks et al. 2000).

**Preplant Burndown Herbicide Use.** Between 20 and 76% of growers used a preplant burndown application (Table 1). Similar to fall herbicide use, the most frequently used herbicides for spring preplant burndown applications across all crop rotations were glyphosate and 2,4-D (Table 3). Furthermore, the use of glyphosate was often four to six times more frequent than 2,4-D, depending on the specific crop rotation. The most frequently used crop-specific herbicides were atrazine and acetochlor in rotations containing corn. In these rotations, glyphosate and 2,4-D were used in preplant burndown applications. A higher percentage of growers in a crop rotation that included GR cotton or GR soybean used glyphosate in their preplant burndown herbicide applications, particularly the growers in the GR cotton / GR soybean rotation. Glyphosate and glyphosate / 2,4-D combinations are effective herbicides for controlling winter annual weeds, and

Table 3. Herbicide applied as preplant burndown for each crop rotation. Data expressed as percentages of producers in each crop rotation who applied each herbicide.

Crop rotation	Atrazine	Acetochlor	Pendimethalin	Isoxaflutole	Glyphosate	2,4-D	Paraquat	
	% of growers making each application							
Continuous GR <sup>a</sup> corn ( <i>n</i> = 23)	22	17	—	—	52	13	—	—
Continuous GR soybean ( <i>n</i> = 183)	—	—	1	—	76	20	—	—
Continuous GR cotton ( <i>n</i> = 74)	—	—	5	—	90	15	—	—
GR corn / GR soybean								
GR corn ( <i>n</i> = 147)	15	13	—	—	40	16	5	—
GR soybean ( <i>n</i> = 155)	—	—	—	—	63	16	2	—
GR cotton / GR soybean								
GR cotton ( <i>n</i> = 29)	—	—	—	—	86	—	—	—
GR soybean ( <i>n</i> = 24)	—	—	—	—	92	—	—	—
GR soybean / non-GR crop								
GR soybean ( <i>n</i> = 41)	—	—	—	—	69	20	1	—
Non-GR crop ( <i>n</i> = 31)	8	—	—	5	48	21	8	—
GR corn / non-GR crop								
GR corn ( <i>n</i> = 21)	10	10	—	—	14	24	—	—
Non-GR crop ( <i>n</i> = 17)	—	—	—	—	59	12	—	—

<sup>a</sup> Abbreviations: GR, glyphosate-resistant.

Table 4. Non-glyphosate herbicides applied for each crop rotation. Data expressed as percentages of producers in each crop rotation who applied each herbicide.

Crop rotation	Non-glyphosate herbicides applied
	% of growers making each application
Continuous GR corn ( <i>n</i> = 36)	47% atrazine 25% acetochlor 8% simazine 6% <i>s</i> -metolachlor 26% chlorimuron 13% flumiclorac 9% 2,4-D 2% <i>s</i> -metolachlor 2% pendimethalin 2% flumioxazin 27% diuron 19% pyriproxyfen 15% MSMA 15% trifloxyulfuron 12% prometryn 11% pendimethalin 8% flumioxazin 8% fluometuron 8% <i>s</i> -metolachlor 2% 2,4-D
Continuous GR soybean ( <i>n</i> = 46)	33% atrazine 28% acetochlor 7% <i>s</i> -metolachlor 5% 2,4-D 15% pendimethalin 11% imazethapyr 9% chlorimuron 8% <i>s</i> -metolachlor 6% acetochlor 6% clethodim 5% 2,4-D 5% flumioxazin
Continuous GR cotton ( <i>n</i> = 62)	20% MSMA 20% <i>s</i> -metolachlor 20% trifloxyulfuron 10% prometryn 15% flumioxazin 10% fluometuron
GR corn / GR soybean GR corn ( <i>n</i> = 181)	—
GR soybean ( <i>n</i> = 67)	15% pendimethalin 11% 2,4-D 9% trifluralin 7% cloransulam 6% imazethapyr 5% flumiclorac 1% acetochlor 1% nicosulfuron 1% <i>s</i> -metolachlor 20% atrazine 20% <i>s</i> -metolachlor 13% mesotriione 12% acetochlor 7% 2,4-D 7% isoxaflutole 6% cipropyralid 6% nicosulfuron 2% pendimethalin 1% trifluralin < 1% cloransulam
GR Cotton / GR Soybean GR cotton ( <i>n</i> = 20)	—
GR soybean ( <i>n</i> = 0) GR soybean / non-GR crop GR soybean ( <i>n</i> = 94)	—
Non-GR crop ( <i>n</i> = 384)	32% atrazine 28% acetochlor 13% <i>s</i> -metolachlor 4% mesotriione 2% glufosinate 16% <i>s</i> -metolachlor 11% 2,4-D 10% acetochlor 9% atrazine 9% mesotriione 5% glufosinate

Table 4. Continued.

Crop rotation	Non-glyphosate herbicides applied
GR corn / non-GR crop GR corn ( <i>n</i> = 47)	32% atrazine 28% acetochlor 13% <i>s</i> -metolachlor 4% mesotriione 2% glufosinate 16% <i>s</i> -metolachlor 11% 2,4-D 10% acetochlor 9% atrazine 9% mesotriione 5% glufosinate
Non-GR crop ( <i>n</i> = 57)	—

the herbicides' relatively low cost makes them attractive options for growers. The usage of glyphosate and 2,4-D is slightly lower for rotations including GR corn, suggesting the utilization of other herbicides. The data in Table 3 support this, showing that herbicides such as atrazine and acetochlor were used in rotations that included GR corn. Johnson et al. (2000) also found that by using glyphosate along with reduced rates of chloroacetamide (acetochlor) or triazine (atrazine) herbicides provided better control of weed species than full rates of chloroacetamide or triazine herbicides without the addition of glyphosate.

**Postemergence Glyphosate Use.** Most growers applied two or fewer postemergence applications of glyphosate during crop growth (Table 1). However, in crop rotations that include GR cotton, 30 to 40% of the growers made three applications of glyphosate. Prior to GR cotton, preemergence and postemergence-directed herbicide applications or cultivation were used to control weeds in cotton (Culpepper and York 1998; Snipes and Mueller 1992a, b; Wilcut et al. 1995). Since the commercialization of GR cotton, more and more growers have moved toward total postemergence weed control programs. Reasons for this change include the lack of herbicides labeled for preplant or preemergence use, the difficulty in obtaining adequate height differential between crops and weeds for postemergence-directed applications, marginal crop tolerance to many of these herbicides, and the specialized equipment needed to make these applications (Askew and Wilcut 1999; Culpepper and York 1999; Snipes and Mueller 1992a, b; Wilcut et al. 1997). The main drawback to a total postemergence program using glyphosate is the lack of residual weed control from glyphosate. Multiple applications of glyphosate to the cotton crop are needed to obtain satisfactory weed control if no other weed control tactics are used.

Two or more postemergence applications of glyphosate in GR soybean were used by 66 to 74% of the growers in a continuous GR soybean or GR cotton / GR soybean cropping system (Table 1). However, only 47 to 50% of the growers used two or more postemergence applications of glyphosate in a GR corn / GR soybean or GR soybean / non-GR crop rotation. Of the growers in continuous GR soybean production, 62% required at least two postemergence applications of glyphosate (Table 1). Of the growers that had GR in their crop rotation, 43 to 53% of them made at

least two applications of glyphosate. The tendency to use fewer postemergence applications of glyphosate may be a function of the soybean row spacing, planting date, or geography (soybean maturity length, duration of crop growth). Soybean weed control programs were dominated by imidazolinones and dinitroaniline herbicides from 1992 to 1996, prior to the introduction of GR soybean. With the introduction of GR soybean, many producers began to rely exclusively on glyphosate for weed management (Young 2006). Another reason for the heavy use of glyphosate in GR soybean is that it fills in the gaps left by many conventional soybean weed management programs (Gianessi 2005).

In GR corn, 31 to 44% of the growers used two postemergence applications of glyphosate, which is relatively less than the frequency of glyphosate use in GR soybean or GR cotton (Table 1). The historical availability of cost-effective non-glyphosate products in corn may partially explain the difference in glyphosate use between crops. For example, atrazine in combination with *s*-metolachlor provides, in most cases, economical, season-long weed control (Thomas et al. 2007). Another reason is the rate of GR corn adoption has been slower than the rate of GR soybean or GR cotton (Johnson and Gibson 2006). The GR trait until recently has not been available in many of the most popular corn hybrids. Glyphosate applications in GR corn can only be made up until the V8 crop stage, or until the crop reaches 76 cm in height (Anonymous 2007). For GR soybean, glyphosate applications can be made up until flowering (R2 stage) (Anonymous 2007). This narrow application window for GR corn may also be a contributing factor to the low adoption of GR corn. Gianessi (2005) found that most corn growers who have adopted GR corn technology have done so because they have difficult-to-control weed problems that necessitate more costly herbicide programs. The work of Johnson et al. (2000) found that the use of glyphosate and atrazine or acetochlor provides better control than the use of glyphosate alone.

**Non-Glyphosate Herbicide Use.** Growers more frequently utilized a non-glyphosate herbicide prior to planting (12 to 18%) and at planting (16 to 25%) in the production of GR corn (Table 1). Corn producers still rely on soil-applied herbicides such as atrazine as the foundation of their weed control programs. Reasons for this are discussed above.

A lower percentage of growers applied non-glyphosate herbicides prior to rotations that included GR soybean or GR cotton (3 to 11%). Common herbicides used prior to planting included diuron, fluometuron, pendimethalin, *s*-metolachlor, and trifluralin. These were commonly used herbicides in weed management programs prior to the development of GR cotton and GR soybean (Young 2006).

During the postemergence timing, rotations that included GR cotton and non-GR crops had 53 to 69% using non-glyphosate herbicides. The herbicides prometryn, pyrithiobac-sodium, MSMA, and trifloxsulfuron are still utilized in cotton for over-the-top and layby applications to achieve satisfactory weed control (Table 4). However only prometryn, pyrithiobac, MSMA, and trifloxsulfuron are commonly used. The herbicide use pattern may change with the release of new GR cotton cultivars in 2006 that allow for later postemergence applications of glyphosate (Huff et al. 2007). Prior to

the release of the enhanced GR cotton trait, glyphosate applications were limited to the four-leaf stage in cotton. Applications later than this could result in fruit abortion and yield reduction (Viator et al. 2003, 2004). With the introduction of enhanced GR cotton in 2006, glyphosate applications are possible from crop emergence until 7 d prior to harvest (Anonymous 2007).

The non-GR crops in the crop rotations included conventional corn, soybean, and rice (*Oryza sativa* L.). For these crops, traditional postemergence weed management practices, such as those herbicides listed in Table 4, are used to achieve acceptable weed control.

During the 2005 growing season, 79 and 85% of the producers in GR soybean did not apply a non-glyphosate herbicide (Table 1). These results are in agreement with the findings of Shaner (2000), who found a decrease in the use of ALS inhibitors, acetyl CoA carboxylase (AACase) inhibitors and protoporphyrinogen oxidase (PPO) inhibitors in soybean since 1993. Gianessi (2005) found that one glyphosate application in some cases substituted for three to four herbicides, often applied separately, with the potential need for tillage to obtain adequate weed control. Glyphosate-based weed control programs are inexpensive, convenient and, given the market value of soybean over the past couple of years, a very attractive option for producers.

Results from this survey show that in most instances non-glyphosate-herbicide based weed management programs have been (GR cotton and GR soybean) or are in the process of being (GR corn) replaced with glyphosate as the core, or sole, herbicide. The longer a GR crop is available to producers, and as GR technology develops and advances, these glyphosate-based weed management programs become more attractive to producers. This trend has been especially evident in GR soybean, which has been available for 11 yr, and for which herbicide-use patterns have progressively moved toward intensive glyphosate programs. Now that new GR technology for cotton is available, and allowing for later applications of glyphosate, one can deduce that this trend will become apparent in GR cotton production as well. The same might be said of GR corn production as the technology matures. Adoption of GR corn in the United States has been slower, due again to several factors. Excellent efficacy of existing herbicide programs, as discussed before, may be a contributing factor. Another factor is that GR corn varieties have not been approved for import into Europe (Gianessi 2005). There has been limited information on the efficacy and economics of GR corn (Thomas et al. 2004a), although current research is addressing this deficit (Gianessi 2005; Johnson et al. 2000; Thomas et al. 2004a, b). The increased interest of domestic ethanol production may address export concerns as more corn is used for ethanol production in the United States.

Researchers have also begun to study the possible adverse effects of weed management systems relying exclusively on glyphosate. Weed shifts and acceleration of glyphosate resistance in weeds are some of the top concerns with these systems (Duke 2005; Shaner 2000; Young 2006). Due to concerns about glyphosate resistance, as well as a number of other management and economic factors, anecdotal data indicate there may be shifts toward greater utilization of soil-

applied herbicides. Thus, a follow-up survey will be of great interest to determine why any shifts in herbicide use patterns may continue to occur.

Grower surveys are a valuable tool for documenting herbicide use patterns and the grower attitudes and perceptions driving decisions regarding herbicide selection. The data from this survey will be invaluable reference material for weed scientists and agricultural analysts in understanding the level of glyphosate herbicide usage, the other primary herbicide tools being utilized, and the current benchmarks for herbicide usage in GR crops. As changes continue to occur in herbicide programs, these data will serve as an important snapshot in time for future reference.

### Acknowledgment

This research was funded by Monsanto Agricultural Products Company. Manuscript approved for publication as Journal Article No. J-11358 of the Mississippi Agricultural and Forestry Experiment Station, Mississippi State University.

### Literature Cited

- Anonymous. 2007. Roundup WEATHERMAX® herbicide product label. St. Louis, MO: Monsanto, [http://www.monsanto.com/monsanto/ag\\_products/pdf/labels\\_msds/roundup\\_weathermax\\_label.pdf](http://www.monsanto.com/monsanto/ag_products/pdf/labels_msds/roundup_weathermax_label.pdf).
- Askew, S. D. and J. W. Wilcut. 1999. Cost and weed management with herbicide programs in glyphosate-resistant cotton (*Gossypium hirsutum*). *Weed Technol.* 13:308–313.
- Ateh, C. M. and R. G. Harvey. 1999. Annual weed control by glyphosate in glyphosate-resistant soybean (*Glycine max*). *Weed Technol.* 13:394–398.
- Barrentine, W. L., E. E. Hartwig, C. J. Edwards Jr., and T. C. Kilen. 1982. Tolerance of three soybean (*Glycine max*) cultivars to metribuzin. *Weed Sci.* 30:344–348.
- Bradley, K. W., E. S. Hagood Jr., and P. H. Davis. 2004. Trumpetcreeper (*Campsis radicans*) control in double-crop glyphosate-resistant soybean with glyphosate and conventional herbicide systems. *Weed Technol.* 18:298–303.
- Burnside, O. C. 1996. The history of 2,4-D and its impact on development of the discipline of weed science in the United States. Pages 5–15 in O. C. Burnside, ed. *Biologic and Economic Assessment of Benefits From Use of Phenoxy Herbicides in the United States*. NAPIAP Report Number 1-PA-96. Washington, DC: USDA.
- Charles, G. W. 1991. A grower survey of weeds and herbicide use in the New South Wales cotton industry. *Aust. J. Exp. Agric.* 31:387–392.
- Corbett, J. L., S. D. Askew, W. E. Thomas, and J. W. Wilcut. 2004. Weed efficacy evaluations of bromoxynil, glufosinate, glyphosate, pyrithiobac, and sulfosate. *Weed Technol.* 18:443–453.
- Culpepper, A. S. and A. C. York. 1998. Weed management in glyphosate tolerant cotton. *J. Cotton Sci.* 4:174–185.
- Culpepper, A. S. and A. C. York. 1999. Weed management and net returns with transgenic, herbicide-resistant, and nontransgenic cotton (*Gossypium hirsutum*). *Weed Technol.* 13:411–420.
- Dillard, H. R., T. J. Wicks, and B. Philp. 1993. A grower survey of diseases, invertebrate pests, and pesticide use on potatoes grown in South Australia. *Aust. J. Exp. Agric.* 33:653–661.
- Duke, S. O. 2005. Taking stock of herbicide-resistant crops ten years after introduction. *Pest Manag. Sci.* 61:211–218.
- Faircloth, W. H., M. G. Patterson, C. D. Monks, and W. R. Goodman. 2001. Weed management programs for glyphosate-tolerant cotton (*Gossypium hirsutum*). *Weed Technol.* 15:544–551.
- Gianessi, L. P. 2005. Economic and herbicide use impacts of glyphosate-resistant crops. *Pest Manag. Sci.* 61:241–245.
- Gibson, K. D., D. E. Hillger, and W. G. Johnson. 2006. Farmer perceptions of weed problems in corn and soybean rotation systems. *Weed Technol.* 20:751–755.
- Gibson, K. D., W. G. Johnson, and D. E. Hillger. 2005. Farmer perceptions of problematic corn and soybean weeds in Indiana. *Weed Technol.* 19:1065–1070.
- Huff, J. A., J. T. Irby, D. M. Dodds, M. T. Kirkpatrick, and D. B. Reynolds. 2007. Glyphosate tolerance in Roundup Ready Flex cotton. *Proc. South. Weed Sci. Soc.* 60:68.
- Johnson, W. G., P. R. Bradley, S. E. Hart, M. L. Buesinger, and R. E. Massey. 2000. Efficacy and economics of weed management in glyphosate-resistant corn (*Zea mays*). *Weed Technol.* 14:57–65.
- Johnson, W. G. and K. D. Gibson. 2006. Glyphosate-resistant weeds and resistance management strategies: an Indiana grower perspective. *Weed Technol.* 20:768.
- Llewellyn, R. S., R. K. Lindner, D. J. Pannell, and S. B. Powles. 2002. Resistance and the herbicide resource: perceptions of Western Australian grain growers. *J. Crop Prot.* 21:1067–1075.
- Padgette, S. R., K. H. Kolacz, X. Delannay, D. B. Re, B. J. LaVallee, C. N. Tinius, W. K. Rhodes, Y. I. Otero, G. F. Barry, and D. A. Eichholz. 1995. Development, identification, and characterization of a glyphosate-tolerant soybean line. *Crop Sci.* 35:1451–1461.
- Reddy, K. N. and K. Whiting. 2000. Weed control and economic comparisons of glyphosate-resistant, sulfonylurea-tolerant, and conventional soybean (*Glycine max*) systems. *Weed Technol.* 14:204–211.
- Shaner, D. L. 2000. The impact of glyphosate-tolerant crops on the use of other herbicides and on resistance management. *Pest Manag. Sci.* 56:320–326.
- Shaw, D. R., W. A. Givens, L. A. Farno, P. D. Gerard, J. W. Wilcut, W. G. Johnson, S. C. Weller, B. G. Young, R. G. Wilson, and M.D.K. Owen. 2009. Using a grower survey to assess the benefits and challenges of glyphosate-resistant cropping systems for weed management in U.S. corn, cotton, and soybean. *Weed Technol.* 23:134–149.
- Snipes, C. E. and T. C. Mueller. 1992a. Influence of fluometuron and MSMA on cotton yield and fruiting characteristics. *Weed Sci.* 42:210–215.
- Snipes, C. E. and T. C. Mueller. 1992b. Cotton (*Gossypium hirsutum*) yield response to mechanical and chemical weed control systems. *Weed Sci.* 42:249–254.
- Snyder, R. L., M. A. Plas, and J. I. Grieshop. 1996. Irrigation methods used in California: grower survey. *J. Irrig. Drain. Eng.* 122:259–262.
- Thomas, W. E., I. C. Burke, and J. W. Wilcut. 2004a. Weed management in glyphosate-resistant corn with glyphosate and halosulfuron. *Weed Technol.* 18:1049–1057.
- Thomas, W. E., I. C. Burke, and J. W. Wilcut. 2004b. Weed management in glyphosate-resistant corn with glyphosate, halosulfuron, and mesotrione. *Weed Technol.* 18:826–834.
- Thomas, W. E., W. J. Everman, J. Allen, J. Collins, and J. W. Wilcut. 2007. Economic assessment of weed management systems in glufosinate-resistant, glyphosate-resistant, imidazolinone-tolerant, and nontransgenic corn. *Weed Technol.* 21:191–198.
- [USDA-NASS] U.S. Department of Agriculture–National Agricultural Statistics Service. 2007. Acreage statistics. Washington, D.C.: USDA National Agricultural Statistics Service, <http://usda.mannlib.cornell.edu/usda/current/Acre/Acre-06-29-2007.pdf>.
- Viator, R. P., P. H. Jost, S. A. Senseman, and J. T. Cothren. 2004. Effect of glyphosate application timings and methods on glyphosate-resistant cotton. *Weed Sci.* 52:147–151.
- Viator, R. P., S. A. Senseman, and J. T. Cothren. 2003. Boll abscission responses of glyphosate-resistant cotton (*Gossypium hirsutum* L.) to glyphosate. *Weed Technol.* 17:571–575.
- Wicks, G. A., G. W. Mahnken, and G. E. Hanson. 2000. Effect of herbicides applied in winter wheat (*Triticum aestivum*) stubble on weed management in corn (*Zea mays*). *Weed Technol.* 14:705–712.
- Wilcut, J. W., D. L. Jordan, W. K. Vencill, and J. S. Richburg III. 1997. Weed management in cotton (*Gossypium hirsutum*) with soil-applied and post-directed herbicides. *Weed Technol.* 11:221–226.
- Wilcut, J. W., A. C. York, and D. L. Jordan. 1995. Weed management programs for oil seed crops. Pages 343–400 in A. E. Smith, ed. *Handbook of Weed Management Programs*. New York: Marcel-Dekker.
- Woodburn, A. T. 2000. Glyphosate: production, pricing and use worldwide. *Pest Manag. Sci.* 56:309–312.
- Young, B. G. 2006. Changes in herbicide use patterns and production practices resulting from glyphosate-resistant. *Weed Technol.* 20:301–307.

*Received March 5, 2008, and approved July 21, 2008.*