**ORGAN TOXICITY AND MECHANISMS** 



# Pre- and postnatal exposure to glyphosate-based herbicide causes behavioral and cognitive impairments in adult mice: evidence of cortical ad hippocampal dysfunction

Yassine Ait-Bali<sup>1,2</sup> · Saadia Ba-M'hamed<sup>1</sup> · Giovanna Gambarotta<sup>3</sup> · Marco Sassoè-Pognetto<sup>2,4</sup> · Maurizio Giustetto<sup>2,4</sup> · Mohamed Bennis<sup>1</sup>

Received: 17 December 2019 / Accepted: 11 February 2020 © Springer-Verlag GmbH Germany, part of Springer Nature 2020

# Abstract

Glyphosate-based herbicides (GBH) are the most widely used pesticides worldwide. Despite considerable progress in describing the neurotoxic potential of GBH, the harmful effects on brain cytoarchitecture and behavior are still unclear. Here, we addressed the developmental impact of GBH by exposing female mice to 250 or 500 mg/kg doses of GBH during both pregnancy and lactation and then examined the downstream effects at the behavioral, neurochemical and molecular levels. We show that pre- and neonatal exposure to GBH impairs fertility and reproduction parameters as well as maternal behavior of exposed mothers. In offspring, GBH was responsible for a global delay in innate reflexes and a deficit in motor development. At the adult age, exposed animals showed a decrease of locomotor activity, sociability, learning and short- and long-term memory associated with alterations of cholinergic and dopaminergic systems. Furthermore, GBH-activated microglia and astrocytes, sign of neuroinflammation event in the medial prefrontal cortex and hippocampus. At the molecular level, a downregulation of brain-derived neurotrophic factor (BDNF) expression and an up-regulation of tyrosine-related kinase receptor (TrkB), NR1 subunit of NMDA receptor as well as tumor necrosis factor  $\alpha$  (TNF $\alpha$ ) were found in the brain of GBH-exposed mice. The present work demonstrates that GBH induces numerous behavioral and cognitive abnormalities closely associated with significant histological, neurochemical and molecular impairments. It also raises fundamental concerns about the ability of current safety testing to assess risks of pesticide exposure during developmental periods of central nervous system.

Keywords Glyphosate · Behavior · Cognition · Acetylcholinesterase · Neuroinflammation · BDNF signaling

**Electronic supplementary material** The online version of this article (https://doi.org/10.1007/s00204-020-02677-7) contains supplementary material, which is available to authorized users.

Mohamed Bennis mbennis@uca.ma

- <sup>1</sup> Laboratory of Pharmacology, Neurobiology and Behavior, Faculty of Sciences Semlalia, Cadi Ayyad University, Bd. Prince My Abdallah, BP. 2390, 40000 Marrakech, Morocco
- <sup>2</sup> Department of Neuroscience, University of Turin, Turin, Italy
- <sup>3</sup> Department of Clinical and Biological Sciences, University of Turin, Turin, Italy
- <sup>4</sup> National Institute of Neuroscience-Italy, Turin, Italy

# Introduction

The American National Academy of Sciences estimated that 3% of brain developmental disorders, such as autism spectrum disorders and learning disabilities, may be directly linked to exposure to environmental chemicals (National Research Council 2000). Existing evidence supports the idea that lifelong susceptibility to anxiety disorders can be partly determined by environmental factors during early development (Gross and Hen 2004). Furthermore, it is important to note that early perturbations of brain development may lead to neurobehavioral disturbances expressed either in childhood or with delayed onset in adulthood (Olney 2002). In line with this, the potential of in utero or early postnatal pesticide exposure to affect brain development has been shown and an emerging literature provides evidence for neurobehavioral consequences resulting from early exposure to organophosphates (OPs) pesticides (for review, see Heyer and Meredith 2017).

Glyphosate (Gly) (N-[phosphonomethyl]glycine; CAS registry number 1071-83-6), the active ingredient present in Roundup® (Monsanto Company, St. Louis, MO), is the most heavily OP herbicide used worldwide (Powles et al. 1997). The herbicidal action of Gly is due to inhibition of a key plant enzyme (5-enolpyruvyl shikimate-3-phosphate synthase) (Franz et al. 1997). Since this enzyme is not present in vertebrates, it has long been assumed that Gly does not affect non-target species. In addition, there are experimental data which supported the safety hypothesis of Gly by showing, on one hand that the blood concentration of Gly reaches its peak approximately 1-2 h after ingestion, and on the other hand, that Gly is poorly metabolized and generally eliminated in the urine and faeces (Chan and Mahler 1992; Brewster et al. 1991). However, over the last few years, several concerns have been raised by the scientific community and regulatory agencies regarding the potential adverse effects of Gly and its adjuvants on the environment and human health (Solomon et al. 2007). Evidence of exposure to Gly has been revealed by its detection in urine samples of people living in farm and non-farm household (Conrad et al. 2017) and in maternal and umbilical cord serum of pregnant women in Thailand (Kongtip et al. 2017). Clinical reports of intoxication with commercial formulations of Gly described negative effects on the nervous system, including Parkinsonism (Barbosa et al. 2001; Wang et al. 2011), anxiety and short-term memory impairments (Nishiyori et al. 2014). Moreover, an epidemiological study highlighted a strong correlation between the increasing application of Gly in agriculture and the occurrence of several neurological diseases, including autism, dementia and anxiety disorder at different ages (Seneff et al. 2015). Concomitantly, experimental studies also revealed neurotoxic effects of glyphosate-based herbicide (GBH). Indeed, oral administration of GBH to pregnant rats alters the activity of brain enzymes in both mothers and offspring (Daruich et al. 2001). These data suggest that GBH could impact maternal behavior and subsequently, offspring's sociability given the crucial role of early maternal care on offspring's abilities (Branchi et al. 2013). However, there have been no systematic studies evaluating the effects of GBH on maternal care and social behavior of their offspring. At the behavioral level, it has been shown that early exposure to GBH results later in life in depressive-like behavior (Cattani et al. 2017) and contradictory effects on the anxiety-related behavior (Gallegos et al. 2016; Baier et al. 2017). Recently, we found that GBH induces cognitive alterations in young male mice, supporting the idea that early exposure to GBH can interfere with brain structures involved in learning and memory leading to cognitive deficits (Bali et al. 2019). However, the effect of prenatal and neonatal exposure to GBH on learning and memory is still unexplored. In this context, the present study was conducted using a multifaceted behavioral battery to assess the profile of gestational and lactational GBH effects in a mouse model. Behavioral assays covering neonatal age and adulthood were selected to measure a range of early reflex development as well as locomotor, affective, sociability and cognitive functions. Mechanistic, cytoarchitectural, neurochemical and molecular mechanisms underlying of GBH-induced neurobehavioral deficits were evaluated in this follow-up study, with the goal to provide benchmark data for GBH risk assessment in the brain.

# **Materials and methods**

#### Pesticide

Roundup herbicide (glyphosate concentration: 360 g/l in the form of glyphosate isopropylamine salt 486 g/l), was used in the liquid commercial form supplied by Monsanto Company (St. Louis, MO, USA). The molecular formula is  $C_6H_{17}N_2O_5P$  (molecular weight: 228.183 g/mol; melting point: 200 °C; density: 1.218 g/cm<sup>3</sup>).

#### Animals

Male and female Swiss mice (3 months old) were obtained from the animal husbandry of the Faculty of Sciences, Cadi Ayyad University, Marrakech, Morocco. The animals were housed in Plexiglas cages ( $30 \times 15 \times 12$  cm) under standard conditions of temperature ( $22 \pm 2$  °C) and photoperiod (12 h/12 h) with food and water ad libitum. All procedures were approved by the Council Committee of Research Laboratories of the Faculty of Sciences, Cadi Ayyad University (Marrakech, Morocco) and conducted in accordance with European Council Directive: EU2010/63. All efforts were made to minimize animal suffering.

#### Doses and protocol of exposure

Females were mated with breeding males (two females for one male) over a day, and were examined on the following day by vaginal plug inspection to assess successful mating. If judged copulated, the female was removed from the cage of the male and housed individually. This was considered as the day 0 of gestation (G0).

Female exposure to GBH through oral gavage occurred daily from G0 to postnatal day 21 (P21) (Fig. 1). Three experimental groups were formed, each one including a minimum of six female mice: a group exposed to a lower dose (250 mg/kg) of GBH, a group exposed to a higher dose (500 mg/kg) of GBH, and a control group which received vehicle (tap water). These doses were selected based on the



**G0:** Gestational day 0 **PND:** Postnatal day

Fig. 1 Experimental protocol

no-observed adverse effect level (NOAEL) indications (i.e.: 500 mg/kg/day) (EPA 1993). The GBH doses used in the present study were higher than the GBH levels to which the population is normally exposed (Solomon 2016). However, as in other toxicological studies, exposure to relatively high doses was used to demonstrate a plausible drug-action (see for example Ford et al. 2017).

# Gestation outcomes, maternal behavior and body weight of pups

To detect any signs of poisoning, all pregnant mice were observed daily from the first administration day (G0) until parturition. In addition, several parameters of maternal behavior, fertility and reproduction were evaluated according to Ema et al. (2008).

### Motor and sensory development assessment

The behavioral testing (negative geotaxis, righting reflex, cliff avoidance and rotarod tests) was performed as described by Ait bali et al. (2016). Animals (n = 10 for each group: control, 250 mg/kg and 500 mg/kg; two males from each litter) were tested during morning sessions starting at 9 a.m. The tests for sensorimotor development assessed during the same day were separated by an interval of 30 min.

#### Adult behavior

From P60, behavioral tests were performed to assess locomotor activity (open field, OF), levels of anxiety (OF and elevated plus-maze, EPM), social interaction (three-chambered sociability test, TCS), working memory (Y-maze), recognition memory (novel object recognition test, NOR), and learning and emotional memory performances (passive avoidance test, PA). The behavior of a total number of ten mice for each group was evaluated between 9 a.m. and 13 p.m and recorded with Ethovision XT Noldus 8.5 video tracking program (Noldus Information Technology b.v., Wageningen, The Netherlands), connected to a video camera (JVC, Japan).

#### **Open field**

This test was performed to assess the general locomotor activity (Walsh and Cummins 1976). Activity monitoring was conducted in a square shaped, white arena, measuring  $50 \times 50 \times 50$  cm. Mice were placed individually into the arena and monitored for 20 min. The assessed parameters were the total distance travelled, the velocity and the time spent in the center.

#### **Elevated plus-maze**

The elevated plus-maze is a widely accepted paradigm used to assess anxiety-like behavior in rodents (Pellow et al. 1985). The elevated plus-maze included two opposing open arms (OA) ( $50 \times 5$  cm) and two closed arms (CA) ( $50 \times 5 \times 15$  cm) joining at a square central area ( $5 \times 5$  cm) to form a plus sign. The entire apparatus was elevated to a height of 45 cm above the floor. Each mouse was tested within a 5-min test session. At the beginning, each mouse was placed individually in the central area facing one of the open arms and allowed to freely explore the maze. The time spent in the OA and CA as well as the number of entries into each arm were quantified. An anxiety index (1 - [([open arm time/total time] + [open arm entries/total number of entries])/2]) was determined according to Cohen et al. (2013).

#### Three-chambered sociability test

The social interaction test was run in a three-chambered arena made of clear glass. Retractable doorways, built into the two dividing walls, controlled access to the side chambers. Each of the two outside chambers had an inverted empty wire cup, one housing a male "stimulus" mouse agematched to the "test" mouse, and the other with a plastic object ("plastic mouse"). The test session began with a 5 min habituation session with the test mouse free to explore the entire arena. This mouse was then briefly confined to the center chamber, while the plastic object was placed in the cup on one side and an adult male mouse on the other side. The "stimulus" mouse and the "plastic mouse" sides were alternated, left and right, between tests. Once the stimuli were in position, the two side doors were simultaneously raised and the test mouse could access all three chambers for 5 min. Automatic monitoring recorded and scored the time spent in contact with each wire cup as well as the number of visits. The apparatus was cleaned between tests using a 70% ethanol/water solution.

#### Y-maze

Y-maze was used to assess working memory performances (Hughes 2004). This three-arm apparatus was made of brown wood (60 cm  $\times$  15 cm  $\times$  30 cm), positioned at equal angles of 120°. Mice were placed at the end of one arm and allowed to freely explore the maze over an 8 min session. The series of arm entries were recorded, and alternation was defined as a triplet of explored arms. Alternation was considered as successful when the three arms were different. The percentage of spontaneous alternation was calculated according to the following equation: % alternation = [(number of alternations)/(total arm entries—2)]  $\times$  100 (Chen et al. 2016).

### Novel object recognition

This test was used to evaluate recognition memory. It was based on the natural preference of mice for a new object with respect to a familiar one. The apparatus consisted of an OF made in Plexiglas ( $50 \times 50 \times 50$  cm) containing two identical or different objects according to the phase of the test. The objects to be discriminated were three plastic objects. The task procedure consists of three phases: habituation, training, and retention phase, according to the protocol described by Bevins and Besheer (2006) for a one trial nonmatching to sample learning procedure. The habituation phase with the apparatus was conducted for 10 min without the presence of the objects. The next day, during the training session, two identical objects were placed in the back corner of the box. The experimental mouse was then placed midway at the front of the box and the total time spent exploring the two objects was recorded for 5 min. During the retention session (1 h after), one of the two identical objects was replaced by a new one and the mouse was allowed to explore the different objects for 10 min. The time spent next to each of the two objects (the familiar one and the novel one) was recorded and discrimination between them was calculated using a discrimination index [DI=(novel object exploration time/total exploration time of two objects)  $\times$  100].

#### **Passive avoidance**

The test is based on the association formed between an aversive stimulus (a foot shock) and a specific environmental context. The apparatus consisted of a two-compartment (light–dark) box. The light compartment  $(10 \times 13 \times 15 \text{ cm})$ was illuminated while the dark one  $(10 \times 13 \times 15 \text{ cm})$  was equipped with energized grid floor, separated by a guillotine door. The entrance of animals to the dark box was punished by an electric foot shock (0.2 mA for 1 s duration). 24 h before the training session, the mice were allowed to explore freely the apparatus for 3 min (habituation). On the training day, each mouse was placed in the center of the light compartment facing away from the guillotine door. After 10 s of adaptation, the guillotine door was opened exposing the dark compartment. When the mouse entered the dark box with all four paws, the guillotine door was closed, and the foot shock was delivered. On the test day, the mouse was returned to the illuminated compartment and the procedure was repeated except that no shock was delivered. The test session was carried out 2 h (short-term memory) or 24 h (long-term memory) after the training. Each time, the latency to enter the dark compartment was recorded. Mice whose latency on the training session exceeded 60 s were excluded from the experiment to minimize the deviation of baseline data. If the animal did not enter the dark compartment during the test within 300 s, the trial was stopped and the final score was established as 300 s. After each session, the apparatus was cleaned using 70% ethanol.

# Determination of acetylcholinesterase enzyme activity

After the behavioral analyses, mice were killed by decapitation, their brains immediately removed from the skull and the PFC and hippocampi were dissected for biochemical analyses. Acetylcholinesterase (AChE) activity was determined in tissue homogenates as described previously (Bali et al. 2019). Briefly, acetylthiocholine (ASCh) was used as a substrate of AChE with Thiocholine (SCh) as a reaction product. The activity of AChE was determined according to the colorimetric method of Ellman et al. (1961) based on the reaction of SCh with DTNB (5,5'-Dithiobis (2-nitrobenzoic acid)), which gives a yellow compound (TNB: S Thio-2-*N* nitro-benzoate) absorbing light at 412 nm. The absorbance of the TNB measured with a spectrophotometer is proportional to the enzymatic activity of AChE. The specific activity is calculated and presented as a percentage:

 $SA = \frac{\Delta OD/min \times 1000}{\varepsilon \times weight of brain tissue},$ 

 $\Delta OD/min$  variation of OD per min, [weight of brain tissue] weight of brain tissue of each mouse (g),  $\varepsilon$  molar extinction coefficient of TNB at 412 nm with  $\varepsilon TNB = 13.6 \times 10^{-6} M^{-1} cm$ .

### Tissue sampling and immunofluorescence

Upon conclusion of behavioral testing, control and treated mice were anesthetized with an intraperitoneal injection of urethane 40% (1 g/kg, from Sigma-Aldrich, France) and transcardially perfused with saline solution (0.9%), followed by ice-cold 4% formaldehyde in phosphate-buffered saline (PBS; 0.1 M). The brains were then removed, post-fixed in the same fixative for 12 h and cryoprotected overnight in 30% sucrose. They were then cut on a freezing cryostat (Leica Microsystems, Germany) into 30 µm frontal sections. The sections containing the substantia nigra pars compacta (SNc), the ventral tegmental area (VTA) and the striatum were used for tyrosine hydroxylase (TH) immunofluorescence, while sections containing PFC and dorsal hippocampus stocked for GFAP and Iba-1 immunofluorescence. The regions of interest were determined according to stereotaxic atlas of Paxinos and Franklin (2001). Sections were kept in PBS containing 0.05% Triton X-100 and 10% normal donkey serum (NDS) for 1 h. Thereafter, they were incubated with the appropriate primary antibodies (mouse monoclonal anti-TH 1:1000, Immunostar cat. 22941; rabbit polyclonal anti-GFAP; 1:500, Abcam, Ab7260; rabbit polyclonal anti-Iba-1; 1:1000; Wako, cat. 019-19741) diluted in PBS with 3% NDS and 0.05 Triton X-100 with gentle stirring at 25 °C. The following day, the sections were rinsed and incubated for 2 h with the adequate secondary antibodies (1:1000; Jackson ImmunoResearch, West Grove, PA, USA) at room temperature for 2 h. After PBS rinsing, the sections were mounted on gelatin-coated glass slides and coverslipped with Dako fluorescence mounting medium (Dako Italia, Milan, Italy).

### **Image analysis**

The TH-immunostained sections were used to assess the number of dopaminergic cells in the SNc and VTA (Bregma: -6.30 mm) and dopaminergic fibers in the striatum (Bregma: 0.48 mm), while the GFAP and Iba-1 immunostained sections were used to assess reactive astocytes and microglia, respectively, in the PFC (Bregma: 3.20 mm) and dorsal hippocampus (Bregma: - 3.80 mm). Three mice per group were used for these analyses. For each brain region, three representative Sections (0.5 µm Z-step size) from anterior to posterior were acquired with a laser scanning confocal microscope (LSM5 Pascal; Zeiss, DE, Germany) using either a  $20 \times objective$  (for TH<sup>+</sup>, GFAP<sup>+</sup> and Iba-1<sup>+</sup> cells count) or a  $40 \times$  objective (for measuring the density of TH<sup>+</sup> innervation) with the pinhole was set at 1 Airy unit. TH<sup>+</sup>, GFAP<sup>+</sup>, and Iba-1<sup>+</sup> cells were manually counted using the point tool in ImageJ software (Image processing and analysis in Java, NIH, USA). The ROI Manager tool in ImageJ software was employed to quantify integral optical density of TH, GFAP and Iba-1 expression. All analyses were carried out by an operator blinded to the experimental groups.

# RNA isolation, cDNA preparation, and quantitative real-time PCR

Twenty-four hours following the completion of behavioral and cognitive tests, RNA was extracted from PFC and hippocampus tissues from control and GBH 500 mg/ kg exposed mice (n=3 each). Isolation of the total RNA was carried out using Trizol (Sigma-Aldrich, St. Louis, MO, USA) according to a previously described protocol (Rio et al. 2010). RNA level was quantified by measuring absorbance at 260 and 280 nm. The final RNA concentration and purity were determined using Nanodrop (Thermo Fisher Scientific, Waltham, MA, USA), and the two ratios DO 260/230 and 260/280 were calculated to verify the quality of the RNAs. Retrotranscription of 1 µg total RNA was carried out in a 25 µl reaction volume containing:  $1 \times RT$ -Buffer, 0.1 µg/µl bovine serum albumin (BSA), 0.05% Triton, 1 mM dNTPs, 7.5 µM Random Hexamer Primers, 40 U RIBOlock, and 200 U RevertAid® Reverse Transcriptase (all RT ingredients were provided by Thermo Scientific). The reaction was performed 10 min at 25 °C, 90 min at 42 °C, 15 min at 70 °C. Quantitative real-time PCR (qRT-PCR) was carried out using an ABI Prism 7300 (Applied Biosystems) detection system. Analyses were performed in technical duplicate and biological triplicate. Data from qRT-PCR experiments were analyzed using the delta cycle threshold ( $\Delta Ct$ ) method. The expression levels were normalized to reference gene: TBP (TATA box Binding Protein). For each tissue (PFC, hippocampus), the  $\Delta Ct$  average of control samples was used as calibrator. Primers were designed using Annhyb software (https://www.bioinformatics.org/annhyb/) and were synthesized by Invitrogen. Primer sequences are reported in Supplementary Table 1.

### **Statistical analysis**

Fertility and reproduction parameters as well as maternal behavior were analyzed by one-way ANOVA, while body weight gain and early sensorimotor endpoint results were analyzed using the repeated measure two-way ANOVA (GBH dose and age), followed by a Holm-Sidak's post hoc test for multiple comparisons. The dataset of behavioral tests in adult mice, enzyme activity results and histological assays were compared between different groups (treated and control) and analyzed using one-way ANOVA, followed by a Holm-Sidak's post hoc for multiple comparisons. The biomolecular data were analyzed with t test. The results are presented as mean  $\pm$  SEM, and a value of p < 0.05 was considered statistically significant. All statistical analyses were carried out using the software SigmaPlot 11.0 for Windows and all graphs were generated with Prism 7.0 for Windows (GraphPad software).

# Results

# Gestation outcome and maternal behavior following GBH exposure

Administration of GBH to pregnant females affected fertility and reproduction parameters. Indeed, the fertility rate and the gestational index were lower in treated groups compared to control. Similarly, both the number of litters and the total number of mice per litter were significantly lower in the GBH-exposed groups. Treatment with GBH 500 mg/kg, but not with the lower dosage, also affected retrieving and nesting index (Table 2). In contrast, the statistical analysis did not reveal a significant difference in the gestation length between treated and control groups ( $F_{(2.18)} = 1.08, p > 0.05$ ) (Table 1).

# GBH decreased body weight of pregnant females and their offspring

The two-way repeated measures ANOVA showed a significant effect of treatment ( $F_{(2,19)} = 11.80$ , p < 0.01) and period of treatment ( $F_{(2,19)} = 211.62$ , p < 0.001) on dam weights. However, the interaction between the two factors was not significant ( $F_{(2,19)} = 0.95$ , p > 0.05). The post hoc comparisons showed that body weight gain was significantly reduced following GBH treatment at both 250 and 500 mg/kg with respect to the control (p < 0.05) (Table 2). Likewise, our results indicated a significant effect of treatment and age on pup's body weight ( $F_{(2,19)} = 11.80$ , p < 0.01;  $F_{(2,19)} = 11.80$ , p < 0.01, respectively) as well as for the interaction between the two factors ( $F_{(2,19)} = 11.80$ , p < 0.01). Furthermore, Holm-sidak comparisons revealed a significant reduction

 
 Table 1
 Reproductive findings in mice given GBH during pregnancy and lactation

	Glyphosate-based herbicide					
	$\overline{0 \text{ (control)}}$	250 mg/kg	500 mg/kg			
No. of females copulated	8	10	24			
No. of pregnant females	7	6	6			
Fecundity index (%)	87	60	25			
No. of death during pregnancy	0	0	1			
Gestation length (days)	$19.8 \pm 0.37$	$20 \pm 0.54$	$19.2 \pm 0.2$			
No. of females with live born	7	6	5			
Gestation index (%)	100	100	83			
No. of females with totally litter loss	0	0	1			
No. of litters	7	6	5			
Total no. of pups born	56	40	35			
No. of pups born alive	51	34	33			
No. of dead pups	5	6	2			
Delivery index (%)	100	100	100			
Lactation index (%)	88.2	79.4	69.6			
Nest building index (%)	100	100	80			
Retrieving index (%)	100	100	80			
% of males	37	45	40			

Fecundity Index (%)=(no. of pregnant females/no. of females copulated)  $\times\,100$ 

Values are given as the mean  $\pm$  SD

Gestation index (%)=(no. of females with live pups born/no. of pregnant females)  $\times\,100$ 

Delivery index (%)=(no. of females delivering/no. of pregnant females)  $\times$  100

Lactation index (%)=(no. of living offspring on day 21/no. of offspring born alive)  $\times$  100

Nesting index = (no. of females building theirs nests for a maximum duration of 30 min/no. of females delivering)  $\times$  100

Retrieving index = (no. of females retrieving over pups for a maximum duration of 30 min/no. of females delivering)  $\times 100$ 

% of males = (no. of males/no. of pups)  $\times 100$ 

of body weight gain in offspring delivered from both GBHexposed dams groups p < 0.05) (Table 2).

# GBH delayed developmental skills of preand postnatally exposed offspring

#### **Righting reflex**

We investigated whether pre- and postnatal exposure to GBH caused atypical sensorimotor skills development in offspring. Two-way repeated measure ANOVA showed a significant main effect of treatment ( $F_{(2,23)} = 13.83$ , p < 0.001) and the expected effect of age, indicating a reduction in righting time as the mice developed ( $F_{(2,23)} = 309.10$ , p < 0.001). Notably, GBH treatment delayed the development of the Table 2GBH affects bodyweight gain of mice mothersand their offspring

	Glyphosate-based herbicide		250 vs control		Post hoc 500 vs control		250 vs 500		
	0 (control)	250 mg/kg	500 mg/kg	t	р	t	р	t	р
Pregnar	псу								
G1	$0.78 \pm 0.34$	$-1.22 \pm 0.48$	$-3.66 \pm 0.14$	3.59	ns	7.98	***	1.94	ns
G7	$5.24 \pm 0.55$	$-2.20 \pm 0.66$	$0.24 \pm 0.49$	5.46	*	8.98	***	1.56	ns
G15	$10.38 \pm 0.67$	$8.12 \pm 0.55$	$5.18 \pm 0.56$	4.06	ns	9.34	***	2.34	ns
G19	$16.34 \pm 0.65$	$14.16 \pm 0.49$	$10.48 \pm 0.78$	3.91	ns	10.53	***	2.93	**
Lactatio	on								
L1	$9.22 \pm 0.95$	$5.04 \pm 1.19$	$3.80 \pm 0.76$	4.34	ns	5.63	*	0.98	ns
L7	$10.00 \pm 0.93$	$6.76 \pm 1.28$	$5.52 \pm 1.01$	3.37	ns	4.66	ns	0.33	ns
L15	$11.06 \pm 0.66$	$7.54 \pm 1.31$	$6.30 \pm 0.83$	3.66	ns	4.95	*	2.80	ns
L19	$11.44 \pm 0.83$	$8.44 \pm 1.01$	$7.20 \pm 0.51$	3.12	ns	4.41	ns	0.33	ns
Litter									
P1	$1.31 \pm 0.02$	$1.15\pm0.07$	$1.05 \pm 0.88$	1.02	ns	1.65	ns	0.51	ns
P7	$2.51 \pm 0.03$	$2.08 \pm 0.14$	$1.91 \pm 0.11$	2.68	ns	3.79	ns	0.90	ns
P15	$5.47 \pm 0.23$	$4.06 \pm 0.22$	$3.17 \pm 0.13$	8.92	***	14.52	***	4.59	***
P21	$7.56 \pm 0.18$	$6.57 \pm 0.24$	$6.43 \pm 0.20$	6.23	**	7.10	***	0.71	ns

G gestation day, L lactation day, P postnatal day

\**p* < 0.05; \*\**p* < 0.01; \*\*\**p* < 0.001; ns (no significant) *p* > 0.05

righting reflex as shown by a significant interaction between the two factors ( $F_{(2,23)} = 8.38$ , p < 0.001). Post hoc analysis revealed that mice treated with 250 mg/kg or 500 mg/kg GBH had slower righting reflexes time than the controls only at P5 (250 vs control mg/kg: q = 7.45, p < 0.001; 500 mg/kg vs control: q = 10.32, p < 0.001) (Fig. 2a).

#### **Negative geotaxis**

The two-way ANOVA revealed a significant effect of treatment ( $F_{(2,23)} = 83.84$ , p < 0.001) and the expected maturation effect reflected by the significant effect of age ( $F_{(2,23)} = 59.44$ , p < 0.001). The interaction was also significant ( $F_{(2,23)} = 23.41$ , p < 0.001). Post hoc comparisons confirmed that mice treated with 500 mg/kg had higher negative geotaxis time compared to the controls at P5, 7 and 9 (q = 19.27, p < 0.001; q = 10.45, p < 0.001 and q = 5.97, p < 0.01, respectively) (Fig. 2b).

#### **Cliff avoidance**

A main effect of treatment was found  $(F_{(2,23)} = 4.05, p < 0.05)$ , as well as the expected effect of age  $(F_{(2,23)} = 23.05, p < 0.001)$ . However, the interaction was not significant  $(F_{(2,23)} = 2.40, p > 0.05)$ . Post hoc comparisons confirmed that mice treated with 500 mg/kg of GBH had slower cliff avoidance time than controls only at P 5 (q = 4.84, p < 0.05) (Fig. 2c).

### **Traction test**

One-way ANOVA analysis showed a significant difference between treated and control groups ( $F_{(2,23)} = 21.38$ , p < 0.001). Post hoc analysis revealed that mice treated with 250 mg/kg or 500 mg/kg GBH showed shorter fall-down latencies than the controls at P10 (250 mg/kg vs control: q=2.41, p < 0.05; 500 mg/kg vs control: q=6.47, p < 0.001). The difference between the two doses of GBH was also statistically significant (250 mg/kg vs 500 mg/kg: q=4.06, p < 0.01) (Fig. 2d).

### **Rotarod test**

The two-way ANOVA analysis revealed that the motor coordination was significantly affected by the treatment  $(F_{(2,23)} = 13.61, p < 0.001)$ , as well as by the age of the animals  $(F_{(2,23)} = 64.04, p < 0.001)$ . Likewise, the interaction was significant  $(F_{(2,23)} = 4.76, p < 0.001)$ . Multiple comparisons confirmed that both GBH-treated groups had lower fall-down latency than the control group on P23 (250 mg/kg vs control: q = 4.95, p < 0.05; 500 mg/kg vs control: q = 4.83, p < 0.05; 500 mg/kg vs control: q = 4.83, p < 0.05; 500 mg/kg vs control: q = 5.42, p < 0.01) (Fig. 2e).

#### Effects of GBH on offspring lasting into adulthood

To investigate whether GBH exposure during prenatal and postnatal developmental might translate into long-lasting consequences on adult behavior, we tested the behavioral



Fig. 2 Pre-and postnatal GBH exposure resulted in neurodevelopmental endpoints changes. a Righting reflex test. b Negative geotaxis test. c Cliff avoidance test. d Traction test. e Rotarod test. Results are

presented as mean  $\pm$  SEM. \*p < 0.05; \*\*p < 0.01; \*\*\*p < 0.001. The "\*" refers to 250 mg/kg or 500 mg/kg vs control groups comparison

repertoire of adults' offspring by assaying locomotor activity, anxiety-like phenotype, social interaction as well as different forms of memory.

### **Open field test**

As indicators of locomotor activity and anxiety-like levels, we recorded the total distance traveled over the open field as well as the velocity and the time spent in the central zone of the maze (anxiety index). One-way ANOVA analysis of the total distance traveled and the percentage of the time spent in the central zone revealed significant differences between treated and control groups ( $F_{(2,17)} = 3.76$ , p < 0.05;  $F_{(2,17)} = 50.67$ , p < 0.001, respectively). However, no difference was found between groups for the velocity  $(F_{(2,17)} = 3.76, p > 0.05)$  (Fig. 3c), even though treated mice seemed slower than controls. Multiple comparisons confirmed that the group of 500 mg/kg exhibited significant decrease of distance traveled compared to the control group (t=2.69, p<0.05) (Fig. 3b). Similarly, post hoc analysis revealed that both GBH-treated groups spent significantly less time in the center of the open field compared to the control (250 mg/kg vs control: t = 6.57, p < 0.001; 500 mg/kg vs control: t = 9.88, p < 0.001). In addition, the same analysis revealed a significant difference between treated groups (t = 3.30, p < 0.01) (Fig. 3d).

#### **Elevated plus-maze**

The open field result was confirmed by EPM data. Indeed, one-way ANOVA analysis of the ratio of time spent in the OA and the anxiety index showed a significant difference between treated and control group ( $F_{(2,17)} = 72.38$ , p < 0.001;  $F_{(2,17)} = 18.17$ , p < 0.001, respectively). However, the analysis of the ratio of entries' number in the OA did not show any statistical difference between groups ( $F_{(2,17)} = 1.05$ , p > 0.05) (Fig. 3g). Multiple comparisons confirmed that the ratio of time spent in the OA was significantly lower in treated groups compared to the control group (250 mg/kg vs control: t = 9.29; p < 0.001; 500 mg/kg vs control: t = 11.26; p < 0.001) (Fig. 3f). Moreover, the anxiety index was significantly higher in treated groups (250 mg/kg vs control: t = 4.36; p < 0.001; 500 mg/kg vs control: t = 5.78; p < 0.001) (Fig. 3h).



**Fig. 3** Pre- and postnatal GBH exposure resulted in behavioral alterations in the offspring during adulthood. **a** Recording of the trajectory of in the OF test. **b**–**d** Effect of GBH on locomotor activity and anxiety-like phenotype in the open field test. **e** Recording of the trajectory of mice in the EPM test. **f**–**h** Effect of GBH on anxiety-like pheno-

type in EPM test. Results are presented as mean  $\pm$  SEM. \*p < 0.05; \*\*\*p < 0.001; #p < 0.05; ##p < 0.01. The "\*" refers to 250 mg/kg or 500 mg/kg vs control groups comparison and the "#" refers to the 250 mg/kg vs 500 mg/kg groups comparison. *OA* open arm, *CA* closed arm

#### Three-chambered sociability

This test was used to investigate the voluntary social interaction of animals. The data analysis revealed a significant interaction between the wire cup (holding mouse vs object) and treatment ( $F_{(2,17)} = 73.53$ , p < 0.001). However, there was no significant main effect of both treatment and wire cup ( $F_{(2,17)} = 2.00$ , p > 0.05;  $F_{(2,17)} = 1.10$ , p > 0.05, respectively). Post hoc comparisons revealed that mice treated either with 250 or 500 mg/kg of GBH spent less time with the wire cup holding another conspecific (250 mg/kg vs control: t=7.42, p < 0.001; 500 mg/kg vs control: t=9.42, p < 0.001; 250 mg/kg vs control: t=9.42, p < 0.001; 250 mg/kg vs 500 mg/kg: t=1.99, p < 0.05) and more time with the wire cup holding the object (250 mg/kg vs control: t=4.89, p < 0.001; 500 mg/kg vs control: t=7.06, p < 0.001; 250 mg/kg vs 500 mg/kg: t=2.16, p < 0.05) compared to the controls (Fig. 4b). We also found a significant effect of the wire cup factor on the visit number as well as its interaction with the treatment factor ( $F_{(2,17)}=26.25$ , p < 0.001;  $F_{(2,17)}=17.89$ , p < 0.001, respectively). Multiple comparisons confirmed



**Fig. 4** Pre- and postnatal GBH exposure resulted in sociability and cognitive alterations in the offspring during adulthood. **a** Recording of the trajectory in the TCS test. **b**, **c** Effect of GBH on social interaction in the TCS test. **d** Effect of GBH on short and long-term memory in the PA test. **e** Effect of GBH on working memory in the Y-maze test. **f**, **g** Effect of GBH exposures on recognition memory

in the NOR test. Results are presented as mean  $\pm$  SEM. \*p < 0.05; \*\*p < 0.01; \*\*\*p < 0.001; #p < 0.05; ##p < 0.01. The "\*" refers to 250 mg/kg or 500 mg/kg vs control groups comparison and the "#" refers to the 250 mg/kg vs 500 mg/kg groups comparison. *M* mouse, *O* object

that only mice exposed to 250 mg/kg showed less visit number of wire cup holding another conspecific (t=4.89, p<0.001), while both GBH-exposed groups showed higher visit number of the wire cup holding the object compared to the controls (250 mg/kg vs control: t=4.39, p<0.001; 500 mg/kg vs control: t=6.32, p<0.001) (Fig. 4c).

#### **Passive avoidance**

We further examined mice by step-through avoidance learning, an experimental paradigm used for assessing learning, short- and long-term memory. The change in the latency to enter into the dark compartment was compared among the three treatment groups and was found to be affected by both treatment and time of test  $(F_{(2,17)} = 13.96)$ , p < 0.01;  $F_{(2,17)} = 31.00$ , p < 0.01, respectively). The analysis revealed also a significant interaction between the two factors ( $F_{(2.17)} = 7.46$ , p < 0.05). Post hoc analysis indicated that only the group exposed to 500 mg/kg of GBH showed a significant decrease of latency after 2 h (500 mg/ kg vs control: t = 6.32, p < 0.001), while both treated groups exhibited a significant decrease of latency 24 h of after the electrical shock administration (250 mg/kg vs control: t = 4.39, p < 0.001; 500 mg/kg vs control: t = 6.32, p < 0.001)(Fig. 4d).

#### Y-maze

The effect of GBH on working memory was evaluated by Y-maze task. One-way ANOVA analysis showed a significant difference in spontaneous activity between treated and control groups ( $F_{(2,17)} = 11.50$ , p < 0.001). Post hoc comparisons confirmed that the spontaneous alternation in GBH-treated mice was lower than that in control mice (250 mg/kg vs control: t = 2.14, p < 0.05; 500 mg/kg vs control: t = 4.78, p < 0.001) and this effect was significantly

#### Novel object recognition

This test was used to assess the potential effects of GBH on recognition memory. One-way ANOVA analysis showed a significant difference in both the ratio of time spent beside the new object and the discrimination index ( $F_{(2,17)}=37.70$ , p < 0.001). Multiple comparisons revealed that GBH-exposed mice spent less time exploring the novel object and had low discrimination index compared to the controls (250 mg/kg vs control: t=5.43, p < 0.001; 500 mg/kg vs control: t=8.58, p < 0.001). In addition, the same analysis revealed a significant difference between treated groups (t=3.15, p < 0.01) (Fig. 4f, g).

# Biochemical, histological and molecular changes within brain of GBH-exposed mice

### **GBH effects on AChE activity**

Because the cholinergic system is closely associated with anxious and cognitive functions (Mineur et al. 2013; Coyle et al. 1983), we sought to assess the impact of GBH exposure on AChE in the supernatants of specific brain areas' homogenates. ANOVA analysis showed significant differences between groups in PFC and whole brain ( $F_{(2,2)} = 12.78$ ; p < 0.05;  $F_{(2,2)} = 13.93$ ; p < 0.05, respectively), while no statistical difference was found in the hippocampus ( $F_{(2,2)} = 5.79$ ; p > 0.05) (Fig. 5c). Post hoc comparisons revealed that only the group exposed to 500 mg/kg of GBH showed a significant decrease of AChE activity in the whole brain (500 mg/kg vs control: t = 5.24, p < 0.05) (Fig. 5a), while the activity of this enzyme was significantly decreased in PFC for both doses 250 and 500 mg/kg doses



**Fig.5** Pre- and postnatal GBH exposure resulted in AChE inhibition. **a** Whole brain. **b** PFC. **c** Hippocampus. Results are presented as mean  $\pm$  SEM. \*p < 0.05. The "\*" refers to 250 mg/kg or 500 mg/kg vs control group comparison

(250 mg/kg vs control: t = 4.85, p < 0.05; 500 mg/kg vs control: t = 3.65, p < 0.05) (Fig. 5b).

#### GBH effect on dopaminergic system

A growing number of data correlate movement control abnormalities with dopaminergic systems dysfunction (Rodríguez et al. 2013; Gallo et al. 2015). To examine the impact of GBH exposure on dopaminergic neurons, we evaluated the expression of tyrosine hydroxylase (TH), a key enzyme involved in dopamine synthesis. Indeed, the quantitative analysis of TH-immunolabeled cells number in the SNc, in the VTA and in the striatum indicated a significant difference between control and treated groups ( $F_{(2,17)} = 17.89$ , p < 0.001;  $F_{(2,17)} = 14.78$ , p < 0.01;  $F_{(2,17)} = 8.41$ , p < 0.05, respectively). Post hoc comparisons confirmed a significant reduction in the number of TH-positive cell bodies (TH<sup>+</sup>) in the SNc (250 mg/kg vs control: 3.66, p < 0.05; 500 mg/kg vs control: 4.47, p < 0.01; 500 mg/kg vs control:



**Fig.6** Pre- and postnatal GBH exposure resulted in dopaminergic circuit defects. **a** Photomicrographs of mice brain cross sections showing the tyrosine hydroxylase (TH)-immunoreactive neurons. **b** Count of TH positive cells in the SNc and **c** in the VTA. **d** The density of TH Immunoreactivity in the striatum. Results are presented

as mean  $\pm$  SEM. \*p < 0.05; \*\*p < 0.01; #p < 0.05. The "\*" refers to 250 mg/kg or 500 mg/kg vs control groups comparison and the "#" refers to the 250 mg/kg vs 500 mg/kg groups comparison. *SNc* substantia nigra pars compacta, *VTA* ventral tegmental area

t=4.91, p<0.01) of GBH-exposed animals compared to the controls (Fig. 6c). Moreover, the same analysis showed that 500 mg/kg treated group showed a significant decrease in the integral optical density of the TH<sup>+</sup> fibers immunofluorescence in the striatum compared to the control and 250 mg/kg groups (500 mg/kg vs control: t=4.07, p<0.01; 250 mg/kg vs 500 mg/kg: t=2.46, p<0.05) (Fig. 6d).

#### **GBH** causes neuroinflammation

Clear evidence showed that organophosphate OPs intoxication is associated with inflammatory responses as revealed the activation of both astrocytes and microglial cells (Banks and Lein 2012). In this perspective, we evaluated neuroinflammation by assessing the expression of both GFAP and Iba-1 proteins in astrocytes and microglial cells, respectively, in different regions of the PFC and the dorsal hippocampus. Interestingly, the quantitative analysis of GFAP and Iba-1 immunofluorescence showed a significant difference among groups. The treated groups showed a significant increase in the number, integral optical density and area occupied by GFAP<sup>+</sup> and Iba-1<sup>+</sup> cells in both PFC and hippocampus (Figs. 7, 8 and supplementary Tables 2, 3).



**Fig.7** Pre- and postnatal GBH exposure resulted in reactive astrocytes. **a**, **k** Micrographs showing the expression of GFAP by immunofluorescence in the PFC and hippocampus. **b**-**d** Count of GFAP positive cells in the PFC and **l**-**n** in the hippocampus. **e**-**g** The integral optical density of GFAP positive cells in the PFC and **o**-**q** in the hippocampus. **h**-**j** Area occupied by GFAP positive cells in the PFC

and **r**–**t** in the hippocampus. Results are presented as mean±SEM. \*p < 0.05; \*\*p < 0.01; \*\*\*p < 0.001; #p < 0.01. The "\*" refers to 250 mg/kg or 500 mg/kg vs control groups comparison and the "#" refers to the 250 mg/kg vs 500 mg/kg groups comparison. *ACCx* anterior cingulate cortex, *PrLCx* prelimbic cortex, *ILCx* infralimbic cortex, *DG* dentate gyrus, *CA1* Ammon's horn 1, *CA3* Ammon's horn 3



**Fig. 8** Pre- and postnatal GBH exposure resulted in reactive microglia. **a**, **k** Micrographs showing the expression of Iba-1 by immunofluorescence in the PFC and hippocampus. **b**–**d** Count of Iba-1 positive cells in the PFC and **l**–**n** in the hippocampus. **e**–**g** The integral optical density of Iba-1 positive cells in the PFC and **o**–**q** in the hippocampus. **h**–**j** Area occupied by Iba-1 positive cells in the PFC and **r**–**t** in the hippocampus. Results are presented as mean  $\pm$  SEM.

# GBH affects the expression of genes associated with neuroinflammation, synaptic plasticity and cell survival

Since one major hallmarks of the inflammatory response is the release of cytokines and chemokines from activated glial cells, we measured the mRNA expression of tumor necrosis factor alpha (TNF $\alpha$ ) by quantitative real-time PCR (qRT-PCR). The statistical analysis showed significant increased levels of TNF $\alpha$  mRNA expression in the hippocampus of



\*p < 0.05; \*\*p < 0.01; \*\*\*p < 0.001; #p < 0.05; #p < 0.01. The "\*" refers to 250 mg/kg or 500 mg/kg vs control groups comparison and the "#" refers to the 250 mg/kg vs 500 mg/kg groups comparison. ACCx anterior cingulate cortex, PrLCx prelimbic cortex, ILCx infralimbic cortex, DG dentate gyrus, CA1 Ammon's horn 1, CA3 Ammon's horn 3

exposed mice compared to controls (t = 5.26, p < 0.05), while no statistical difference was found in the PFC (t = 0.41, p > 0.05) (Fig. 9a).

It is well reported that the activation of NMDA glutamate receptors (NMDARs) is crucial for synaptic plasticity underlying both short- and long-term memory storage, and that the disruption of these receptors function can cause profound cognitive deficits (Lewis 1997; Malenka and Nicoll 1999). Moreover, the interaction between proinflammatory cytokines and the glutamatergic



**Fig. 9** Pre- and postnatal GBH exposure resulted in genes expression in the PFC and hippocampus of adult progeny. **a** qRT-PCR analysis of TNF $\alpha$  transcript. **b**-**d** qRT-PCR analysis of NMDA receptor subunits transcripts. **e** qRT-PCR analysis of BDNF and **f** its receptor TrkB

transcripts. For each tissue (PFC, Hippocampus), the gene expression is shown relatively to its control samples. Results are presented as mean  $\pm$  SEM. \*p <0.05; \*\*p <0.01. The "\*" refers to 250 mg/kg or 500 mg/kg vs control groups comparison

neurotransmission has been previously described (Fogal and Hewett 2008). Indeed, excessive activation of NMDA receptors leads to elevated calcium influx into cells, which perturbs mitochondria and increases generation of reactive oxygen species, inflammation leading ultimately to neurodegeneration. To test whether GBH could also impact glutamatergic signaling, we assessed the expression level of genes encoding for different NMDA receptors subunits. Our results showed that GBH induced significant increase in the mRNA expression of NR1 subunit in the PFC (t=4.92, p < 0.01) (Fig. 9b), while no significant differences could be observed for both NR2A and NR2B subunits in either the PFC or the hippocampus (p > 0.05) (Fig. 9c, d).

The role of various memory-associated neurotrophins, such as brain-derived neurotrophic factor (BDNF), has also been suggested in memory dysfunction (Alonso et al. 2005). Based on our data related to cognitive deficits, we assessed the impact of GBH exposure on the expression of brain-derived neurotrophic factor (BDNF) and its receptor tyrosine regulated kinase B (TrkB) within the PFC and hippocampus. The analysis showed a statistically significant decreasing effect of GBH on the expression of BDNF (t=3.76, p<0.05) and a significant increasing effect on the expression of its receptor TrkB (t=3.08, p<0.05) in the cortices of treated mice (Fig. 9e, f). However, no statistical significant

differences were found in the hippocampi (t=1.65, p>0.05; t=0.05, p>0.05, respectively) (Fig. 9e, f).

# Discussion

In the present study, we characterized the long-term effects of GBH exposure, which may interfere with brain development and lead to permanent abnormalities. Thus, we developed an experimental murine model of chronic pre- and postnatal exposure to GBH and examined the downstream effects at the behavioral, histological and molecular levels in pups and adult mice.

#### **GBH** affects reproduction parameters

Our results showed that continuous exposure to GBH during pregnancy and lactation affected fertility and reproduction in mouse females. Furthermore, GBH exposure exerted negative effects also on maternal behavior (Rodier 1995). Reproduction and fertility toxicity observed in this work are in agreement with the results published in a report of the World Health Organization (WHO 1994). Indeed, exposure of 3500 mg/kg of Gly in rats from day 6 to 19 of pregnancy caused increased maternal mortality, augmented incidence of early resorptions, and decreased number of implantations and viable fetuses (WHO 1994). Although the precise reprotoxicity mechanism is yet to be clarified, it has been previously reported that Gly targets two crucial steps of steroidogenesis in mammals: at the first rate-limiting level of mitochondrial cholesterol transport (Walsh et al. 2000), and at the last irreversible conversion of sexual androgens into estrogens, via a direct action on the aromatase enzyme (Richard et al. 2005).

### **GBH delayed sensorimotor development**

So far, it has been shown that GBH is able to cross the placental barrier which could possibly alter the developmental process of the fetus (Mose et al. 2008). More recently, the report published by the non-governmental American organization "Moms Across America" suggested that detection of relatively high levels of Gly in breast milk in three out of ten women raised a bioaccumulation problem (Honeycutt and Rowlands 2014) and, therefore, posed potential risks for breastfed offspring. Our results support these hypotheses: early after birth, we found that GBH-exposed offspring displayed delayed developmental reflexes likely related to abnormal maturity of sensorimotor (Fox 1965; Iezhitsa et al. 2001), vestibular and/or proprioceptive functions (Secher et al. 2006; Santillán et al. 2010). In the present work, we also highlighted that offspring prenatally exposed to GBH shows a significantly shorter latency to fall from the rotarod than controls which could be due to alterations in the cerebellar function, since this center is heavily involved in the rotarod performance (Hamm et al. 1994). The reduction in the motor coordination could also be due to a failure in neuromuscular maturity as mentioned by Perez-Reyes et al. (1998). This suggestion is further reinforced by our results obtained through the suspension test showing reduced muscle strength in exposed animals. In sum, the pattern of developmental deficits emerged from this study is in agreement with previous human and animal studies showing similar reflexes defects (Engel et al. 2011; Laugeray et al. 2014; Lan et al. 2017).

# GBH reduces locomotor activity and affects dopaminergic system

Normal development of neonatal reflexes can be considered as an index of brain maturation, and late acquisition of these milestones represent a predictive factor of other behavioral changes in adulthood. In line with this idea, our results show that adult progeny from GBH-treated groups presented locomotor hypoactivity which is in agreement with Gallegos et al. (2016), showing that rats exposed to 200 mg/kg of GBH during pregnancy and lactation are hypoactive, and with our previous observations in juvenile mice (Ait bali et al. 2017). Albeit locomotor hypoactivity was previously reported following prenatal exposure to GBH (Gallegos et al. (2016), the neuronal basis still unclear. Thus, to generate a mechanistic understanding whereby GBH produces locomotor hypoactivity, we assessed the outcomes of GBH exposure on dopaminergic system. The nigrostriatal pathway, crucial in movement control, has been shown to be vulnerable to herbicides (Thiruchelvam et al. 2000; Rodríguez et al. 2013). Considering the idea that reduction in locomotor activity is positively correlated with loss of dopaminergic neurons in the SNc (Bano et al. 2014; Gallo et al. 2015), one of the most intriguing findings of our work is that GBH-exposed mice show a robust decrease in the number of dopaminergic neurons, both in the SNc and VTA, and fibers, in the striatum. The decrease of dopaminergic cells we observed may be due to apoptotic events triggered by elevated oxidative stress observed after Gly exposure (Astiz et al. 2009). Our results are in agreement with Hernandez-Plata et al. (2015) showing that intraperitoneal exposure to 150 mg/kg of Gly for 2 weeks produces a decrease of DA level in the striatum of adult rats that is associated with hypoactivity. Interestingly, human reports support the central effects of GBH on basal ganglia circuits following intoxication with commercial formulations of Gly, described as alterations in the GP and SNc closely related to Parkinsonian syndrome (Barbosa et al. 2001). Finally, although the hypoactivity induced by Gly administration is similar to that observed after administration of a DA antagonist (Hernandez-Plata et al. 2015), Gly can affect other neurotransmitter systems involved in motor control (Martinez et al. 2018). Quantitative analysis of serotonin, dopamine and norepinephrine levels confirmed a dramatic loss of these neurotransmitters mainly in the striatum, PFC and hippocampus of rats exposed to 35, 75, 150 or 800 mg/kg of Gly (Martinez et al. 2018), strongly indicating that further studies are needed to decipher the contribution of these systems to GBH-produced hypoactivity.

### GBH increase anxiety level and reduces social behavior

Our data clearly showed that early exposure to GBH increases anxiety levels, which is consistent with another study (Baier et al. 2017) in which Gly was administered to adult mice by intranasal irrigation. In contrast, Gallegos et al. (2016) reported that rats exposed to 100 or 200 mg/kg of GBH (Glifloglex<sup>®</sup>) during pregnancy and lactation experienced low levels of anxiety in adulthood. The difference observed between our results and those of Gallegos et al. (2016) could be explained by the diversity of the commercial formula of Gly administered in the two studies—i.e. different chemical composition in terms of adjuvants. Indeed, it has been pointed out that the herbicidal activity of Gly is potentiated by the presence of adjuvants (Mitchell et al. 1987), or to the possible

synergy between Gly and the other formulation ingredients (El-Shenawy 2009; Mesnage et al. 2013), thus supporting their potential role in the differential effects of Gly on anxiety levels.

It is well established that neuroinflammation expressed by overproduction of proinflammatory cytokines and excessive activation of glia plays an important role in the etiology of many psychiatric disorders including anxiety (Holloway-Erickson et al. 2012). Accordingly, the neuroinflammatory condition that we report in GBH-treated mice, associated with increased anxiety levels, supports this idea. The PFC is strongly involved in the expression of behavioral and autonomic responses to emotionally relevant stimuli, and imaging studies highlight abnormalities in the structure and function of this region in patients with mood disorder (Kennedy et al. 2001; Drevets 2001). We hypothesized that GBH-induced neuroinflammation of this cortical area could lead to cellular disorganization within the PFC since we already observed that GBH exposure induced loss of serotoninergic fibers within this structure in young mice (Ait bali et al. 2017). Nevertheless, the outcomes of neuroinflammation within the PFC was not investigated in the present work and future studies will be necessary to completely understand the mechanism whereby GBH-induced neuroinflammation produces functional alteration of the PFC.

Our results show that perinatal exposure to GBH has a strong effect on the social skills of adult mice, as reflected by significant alterations in the three-chamber test, an analysis conventionally used to test sociability also in murine models of autism (Silverman et al. 2010). Despite that mice are a social species, engaging in high levels of social interactions and sexual/parenting behaviors (Arakawa et al. 2008), perinatally GBH-exposed mice are unable to distinguish a social partner from a new object in adulthood, indicating that Gly effects on social behavior are long lasting. Our results are in agreement with those of Laugeray et al. (2014) who found that exposure to another OP, glufosinate, during pregnancy and suckling periods reduced sociability in mice. There is evidence that the early social environment (e.g.: maternal care and interactions) has a profound impact on the offspring ability to develop normal social skills (Branchi et al. 2013). Our results showing that GBH significantly alters maternal weight gain, nest building capacity, and retrieving behavior suggest that atypical sociability observed in GBN-exposed mice could be, at least in part, due to physiological or behavioral deficits induced in the mothers. However, the cellular origins and circuit mechanisms contributing to these phenotypes have yet to be identified, a future goal that will shed light into the pathogenic mechanisms that underlie social interaction impairment after GBH exposure.

# GBH induces cognitive and cholinergic system impairment and triggers neuroinflammation

The current study showed that GBH affects recognition, working and contextual memory. These results are similar to those recently published by our laboratory in juvenile mice exposed to the same doses of GBH (Bali et al. 2019), and support those of Gallegos et al. (2018) revealing a significant impairment of recognition memory in adult rat prenatally exposed to 100 or 200 mg/kg of GBH. These results are in agreement with clinical reports showing that populations accidentally exposed to the Gly have developed short- and long-term memory impairments accompanied with hippocampal lesion (Barbosa et al. 2001; Nishiyori et al. 2014). Human and animal studies have implicated the central cholinergic system as an important regulator of cognitive functions such as learning and memory. Indeed, brain cholinergic hypofunction associated with memory deficits was previously observed in patients with Alzheimer's disease (Coyle et al. 1983). Similarly, learning and cognitive defects paralleled with cholinergic alteration were recorded in different animal models (for review, see Voorhees et al. 2017). In agreement with these studies, our results showed that GBH triggers cholinergic crisis expressed by significant decrease of AChE activity in the PFC and the whole brain. Although Gly is a weak inhibitor of the AChE activity (Larsen et al. 2016), the results of the present work corroborate the previous data from our laboratory showing that GBH reduced AchE activity in young mice (Bali et al. 2019). These findings support the previous evidence showing that the commercial formulation is more toxic than Gly (Folmar et al. 1979; Richard et al. 2005). It is well known that in addition to its involvement in adult neurotransmission by its catalytic functions (hydrolysis of acetylcholine), AChE also plays a morphogenic role during the development of the nervous system (Grisaru et al. 1999). Therefore, inhibition of AChE activity observed following GBH exposure could interfere with the morphogenic role of this enzyme and would be the main mechanism inherent to the neurobehavioral alterations induced by these compounds. Furthermore, the cognitive impairments associated with the AChE inhibition following exposure to GBH could be also explained by over-stimulation and persistent activation of both muscarinic and nicotinic receptors due to the accumulation of acetylcholine in the synaptic cleft (Scheffel et al. 2018). This finding might offer a partial explanation for the complaints on memory loss reported by workers chronically exposed to OPs compounds even though the neurotoxicity of Gly or GBH may not be entirely due to disturbances of the cholinergic system.

It is recognized that the co-operation of the PFC and the hippocampus is vital for fundamental cognitive functions, and that disconnection or damage to either one of these two brain regions induces impaired cognitive behaviors (Yoon et al. 2008). Furthermore, evidence linking inflammation to neurodegeneration and cognitive defects (Dziedzic 2006) suggests the possibility that induction of inflammation by chronic exposure to OPs may be mechanistically related to deficits in cognitive ability. Based on these, our morphological analysis showed an increase in the GFAP expression in the PFC and hippocampus of adult progeny from GBHtreated groups. Likewise, GBH elevated the expression of Iba-1 in PFC and hippocampus indicating that microglia, as the most sensitive immune cells in the brain, was activated after GBH exposure. In addition, our study demonstrated a significant increase in the expression of proinflammatory cytokines TNFa in the PFC and hippocampus after GBH exposure. These results indicate that repeated exposure to GBH can trigger neuroinflammation response in PFC and hippocampus of exposed mice. In agreement with our results, several previous studies elaborated that exposure to OPs induced glial cells activation as well as cytokine and chemokine elevation contributing to neuronal damage in the hippocampus and piriform cortex. Indeed, soman (an OP nerve agent) intoxication upregulates the GFAP expression and activates microglia in many regions of the brain, including the hippocampus (Angoa-Perez et al. 2010). Similarly, chronic administration of the chlorpyrifos increases GFAP expression in rat hippocampus (Lim et al. 2011).

# GBH increase the expression of glutamatergic receptor and affects cell survival pathway

There is evidence that neuroinflammation exacerbates neuronal damage due to excitotoxicity via interactions between proinflammatory cytokines and glutamatergic pathways (Fogal and Hewett 2008). Indeed, excessive activation of NMDA receptors leads to elevated calcium influx into cells, which perturbs mitochondria and increases generation of reactive oxygen species ultimately leading to cells death. Our biomolecular analysis corroborates these findings showing that, in addition to neuroinflammation event, GBH increases the transcript level of NR1 subunit of NMDA receptors in the PFC. Then, it would seem that GBH could facilitates and enhances NMDA activation either through: (1) increase of NMDA receptors expression, (2) excessive release of glutamate by activated astrocytes or (3) reduction of glutamate uptake and metabolism within glial cells, associated with an increased release of this neurotransmitter in the synaptic cleft (Cattani et al. 2014). Taken together, our data present additional mechanistic bases to explain excitotoxic condition associated with cognitive impairments observed following GBH exposure.

In the PFC and hippocampus, signaling through the BDNF pathway plays an important role in the survival, maintenance and growth and promotes neuronal plasticity and neurogenesis, processes that have been described as potential cellular mechanisms for learning and memory (Sakata et al. 2013). Our results showed that mice exposed to GBH display decreased BDNF levels in PFC, which is coupled with impaired cognitive impairments. In the brain, BDNF binds to TrkB receptor causing its phosphorylation and subsequent activation of the intracellular signaling cascade to promote synaptic plasticity (Cunha et al. 2010). Our study demonstrated that the GBH increased TrkB transcript in the PFC. Therefore, the increased TrkB mRNA level we observed in the PFC after exposure to GBH may reflect a compensatory mechanism to balance reduction of BDNF expression. In agreement with our results, Jain et al. (2013) demonstrated that chronic exposure to Trizophos, another OP, significantly reduced the mRNA expression and protein levels of BDNF in rat hippocampi correlated with learning and memory deficits (Jain et al. 2013).

In summary, our results show that GBH has pervasive harmful effects when administered during the highly sensitive pre- and postnatal periods. Our results indicate that GBH exposure induces multiple behavioral abnormalities involving motor, emotional, social and cognitive functions and targets CNS integrity, affecting cholinergic, dopaminergic and glutamatergic systems as well as neuroinflammation and cellular stress induction. These results strongly shed light on additional (noncholinergic) mechanisms mediating neurological injury induced by this OP pesticide. All data presented about the behavioral changes as well as brain abnormalities induced by GBH pave the way for further analyses: brain microarray analysis is needed to gain a better understanding regarding the molecular and cellular mechanisms involved in the neurodevelopmental effects of GBH. Such analysis will, with no doubt, contribute to our knowledge of the constellations of genes involved in pathological processes mediated by pre- and postnatal exposure to GBH.

Acknowledgements The authors are grateful to Prof I. Elatiri for technical assistance and to Prof A. Buffo for fruitful discussions about astrocytes and microglia analysis. Y. A-B. was a recipient of a fellowship from the Italian Ministry of Foreign Affairs and International Cooperation. We are grateful to Fondo di Beneficienza Intesa Sanpaolo (Italy) for financial support.

# References

- Ait Bali Y, Ba-Mhamed S, Bennis M (2017) Behavioral and immunohistochemical study of the effects of subchronic and chronic exposure to glyphosate in mice. Front Behav Neurosci 11:146
- Ait-Bali Y, Ba-Mhamed S, Bennis M (2016) Prenatal Paraquat exposure induces neurobehavioral and cognitive changes in mice offspring. Environ Toxicol Pharmacol 48:53–62
- Alonso M, Bekinschtein P, Cammarota M, Vianna MRM, Izquierdo I, Medina JH (2005) Endogenous BDNF is required for longterm memory formation in the rat parietal cortex. Learn Mem 12:504–510

- Angoa-Pérez M, Kreipke CW, Thomas DM, Van Shura KE, Lyman M, McDonough JH, Kuhn DM (2010) Soman increases neuronal COX-2 levels: possible link between seizures and protracted neuronal damage. Neurotoxicology 31(6):738–746
- Arakawa H, Blanchard DC, Arakawa K, Dunlap C, Blanchard RJ (2008) Scent marking behavior as an odorant communication in mice. Neurosci Biobehav Rev 32(7):1236–1248
- Astiz M, Alaniz MJTd, Marra CA (2009) Effect of pesticides on cell survival in liver and brain rat tissues. Ecotoxicol Environ Saf 72(7):2025–2032
- Baier CJ, Gallegos CE, Raisman-Vozari R, Minetti A (2017) Behavioral impairments following repeated intranasal glyphosatebased herbicide administration in mice. Neurotoxicol Teratol 64:63–72
- Bali YA, Kaikai NE, Ba-Mhamed S, Bennis M (2019) Learning and memory impairments associated to acetylcholinesterase inhibition and oxidative stress following glyphosate based-herbicide exposure in mice. Toxicology 415:18–25
- Banks CN, Lein PJ (2012) A review of experimental evidence linking neurotoxic organophosphorus compounds and inflammation. Neurotoxicology 33(3):575–584
- Bano F, Ahmed A, Parveen T, Haider S (2014) Anxiolytic and hyperlocomotive effects of aqueous extract of Nigella sativa L. seeds in rats Pak. J Pharm Sci 27:1547–1552
- Barbosa ER, Leiros da Costa MD, Bacheschi LA, Scaff M, Leite CC (2001) Parkinsonism after glycine-derivate exposure. Mov Disord 16:565–568
- Bevins RA, Besheer J (2006) Object recognition in rats and mice: a one-trialnon-matching-to sample learning task to study recognition memory. Nat Protoc 1:1306–1311
- Branchi I, Curley JP, Dandrea I, Cirulli F, Champagne FA, Alleva E (2013) Early interactions with mother and peers independently build adult social skills and shape BDNF and oxytocin receptor brain levels. Psychoneuroendocrinology 38:522–532
- Brewster DW, Warren J, Hopkins WE II (1991) Metabolism of glyphosate in Sprague-Dawley rats: tissue distribution, identification, and quantitation of glyphosate-derived materials following a single oral dose. Fundam Appl Toxicol 17(1):43–51
- Cattani D, Cesconetto PA, Tavares MK, Parisotto EB, De Oliveira PA, Rieg CEH, Zamoner A (2017) Developmental exposure to glyphosate-based herbicide and depressive-like behavior in adult offspring: implication of glutamate excitotoxicity and oxidative stress. Toxicology 387:67–80
- Cattani D, de Liz Oliveira Cavalli VL, Heinz Rieg CE, Domingues JT, Dal-Cim T, Tasca CI, Zamoner A (2014) Mechanisms underlying the neurotoxicity induced by glyphosate-based herbicide in immature rat hippocampus: involvement of glutamate excitotoxicity. Toxicology 320:34–45
- Chan P, Mahler J (1992) NTP technical report on the toxicity studies of Glyphosate (CAS No. 1071–83-6) Administered in dosed feed to F344/N rats and B6C3F1 mice. Toxic Rep Ser 16:1-D
- Chen Z, Huang C, Ding W (2016) Z-Guggulsterone improves the scopolamine-induced memory impairments through enhancement of the BDNF signal in C57BL/6J mice. Neurochem Res 41(12):3322–3332
- Cohen H, Matar MA, Joseph Z (2013) Animal Models of post-traumatic stress disorder. Current protocols in neuroscience. John Wiley and Sons (chapter 9, unit 9.45)
- Conrad A, Schroter-Kermani C, Hoppe HW, Ruther M, Pieper S, Kolossa-Gehring M (2017) Glyphosate in German adults—time trend (2001 to 2015) of human exposure to a widely used herbicide. Int J Hyg Environ Health 220:8–16
- Coyle JT, Price DL, Delong MR (1983) Alzheimer's disease: a disorder of cortical cholinergic innervation. Science 219(4589):1184–1190
- Cunha. (2010) A simple role for BDNF in learning and memory? Front Mol Neurosci 3:1

- Daruich J, Zirulnik F, Gimenez MS (2001) Effect of the herbicide glyphosate on enzymatic activity in pregnant rats and their fetuses. Environ Res 85(3):226–231
- Drevets WC (2001) Neuroimaging and neuropathological studies of depression: implications for the cognitive-emotional features of mood disorders. Curr Opin Neurobiol 11:240–249
- Dziedzic T (2006) Systemic inflammatory markers and risk of dementia. Am J Alzheimer's Dis Other Dement 21(4):258–262
- El-Shenawy NS (2009) Oxidative stress responses of rats exposed to roundup and its active ingredient glyphosate. Environ Toxicol Pharmacol 28:379–385
- Ellman GL, Courtney KD, Andres V, Featherstone RM (1961) A new and rapid colorimetric determination of acetylcholinesterase activity. Biochem Pharmacol 7(2):88–95
- Ema M, Hara H, Matsumoto M (2008) Evaluation of developmental neurotoxicity of polysorbate 80 in rats. Reprod Toxicol 25:89–99
- Engel SM, Wetmur J, Chen J, Zhu C, Barr DB, Canfield RL, Wolff MS (2011) Prenatal exposure to organophosphates, paraoxonase 1, and cognitive development in childhood. Environ Health Perspect 119(8):1182–1188
- EPA (1993) Registration eligibility decision (RED) glyphosate. Office of Prevention, Pesticides and Toxic Substances. EPA-738-F-93-011. Washington, DC.
- Fogal B, Hewett SJ (2008) Interleukin-1β: a bridge between inflammation and excitotoxicity? J Neurochem 106(1):1–23
- Folmar LC, Sanders HO, Julin AM (1979) Toxicity of the herbicide glyphosate and several of its formulations to fish and aquatic invertebrates. Arch Environ Contam Toxicol 8(3):269–278
- Ford B, Bateman LA, Gutierrez-Palominos L, Park R, Nomura DK (2017) Mapping proteome-wide targets of glyphosate in mice. Cell Chem Biol 24:133–140
- Fox WM (1965) Reflex and behavioural development of the mouse. Anim Behav 13(234):24
- Franz JE, Mao MK, Sikorski JA (1997) Glyphosate: a unique global herbicide (ACS Monograph No. 189). American Chemical Society, Washington, DC
- Gallegos CE, Baier CJ, Bartos M, Bras C, Domínguez S, Mónaco N, Minetti A (2018) Perinatal glyphosate-based herbicide exposure in rats alters brain antioxidant status, glutamate and acetylcholine metabolism and affects recognition memory. Neurotox Res 34(3):363–374
- Gallegos CE, Bartos M, Bras C, Gumilar F, Antonelli MC, Minetti A (2016) Exposure to a glyphosate-based herbicide during pregnancy and lactation induces neurobehavioral alterations in rat offspring. Neurotoxicology 53:20–28
- Gallo EF, Salling MC, Feng B, Moron JA, Harrison NL, Javitch JA (2015) Upregulation of dopamine D2 receptors in the nucleus accumbens indirect pathway increases locomotion but does not reduce alcohol consumption. Neuropsychopharmacology 40:1609–1618
- Grisaru D, Sternfeld M, Eldor A, Glick D, Soreq H (1999) Structural roles of acetylcholinesterase variants in biology and pathology. Eur J Biochem 264(3):672–686
- Gross C, Hen R (2004) The developmental origins of anxiety. Nat Rev Neurosci 5:545–552
- Hernández-Plata I, Giordano M, Díaz-Muñoz M, Rodríguez VM (2015) The herbicide glyphosate causes behavioral changes and alterations in dopaminergic markers in male Sprague-Dawley rat. Neurotoxicology 46:79–91
- Heyer DB, Meredith RM (2017) Environmental toxicology: sensitive periods of development and neurodevelopmental disorders. Neurotoxicology 58:23–41
- Holloway-Erickson CM, McReynolds JR, McIntyre CK (2012) Memory-enhancing intra basolateral amygdala infusions of clenbuterol increase Arc and CaMKIIalpha protein expression in the rostral anterior cingulate cortex. Front Behav Neurosci 6:17

- Honeycutt Z, Rowlands H (2014) Glyphosate testing report: findings in American mothers' breast milk, urine and water. Unpublished report, dated 7 April 2014, available from the websites of—Moms Across America|| and—Sustainable Pulse
- Hughes RN (2004) The value of spontaneous alternation behavior (SAB) as a test of retention in pharmacological investigations of memory. Neurosci Biobehav Rev 28(5):497–505
- Iezhitsa IN, Spasov AA, Bugaeva LI (2001) Effects of bromantan on offspring maturation and development of reflexes. Neurotoxicol Teratol 23(2):213–222
- Jain S, Banerjee BD, Ahmed RS, Arora VK, Mediratta PK (2013) Possible role of oxidative stress and brain derived neurotrophic factor in triazophos induced cognitive impairment in rats. Neurochem Res 38(10):2136–2147
- Kennedy SH, Evans KR, Kruger S, Mayberg HS, Meyer JH, McCann S (2001) Changes in regional brain glucose metabolism measured with positron emission tomography after paroxetine treatment of major depression. Am J Psychiatry 158:899–905
- Kongtip P, Nankongnab N, Phupancharoensuk R, Palarach C, Sujirarat D, Sangprasert S, Woskie SR (2017) Glyphosate and paraquat in maternal and fetal serums in Thai women. J Agromed 22:282–289
- Lan A, Kalimian M, Amram B, Kofman O (2017) Prenatal chlorpyrifos leads to autism-like deficits in C57B16/J mice. Environ Health 16(1):43
- Larsen KE, Lifschitz AL, Lanusse CE, Virkel GL (2016) The herbicide glyphosate is a weak inhibitor of acetylcholinesterase in rats. Environ Toxicol Pharmacol 45:41–44
- Laugeray A, Herzine A, Perche O, Hébert B, Aguillon-Naury M, Richard O, Mortaud S (2014) Pre- and postnatal exposure to low dose glufosinate ammonium induces autism-like phenotypes in mice. Front Behav Neurosci 8:1–14
- Lewis DA (1997) Development of the prefrontal cortex during adolescence: insights into vulnerable neural circuits in schizophrenia. Neuropsychopharmacology 16:385–398
- Lim KL, Tay A, Nadarajah VD, Mitra NK (2011) The effect of consequent exposure of stress and dermal application of low doses of chlorpyrifos on the expression of glial fibrillary acidic protein in the hippocampus of adult mice. J Occup Med Toxicol 6(1):4
- Malenka RC, Nicoll RA (1999) Long-term potentiationa decade of progress? Science 285:1870–1874
- Martínez MA, Ares I, Rodríguez JL, Martínez M, Martínez-Larrañaga MR, Anadón A (2018) Neurotransmitter changes in rat brain regions following glyphosate exposure. Environ Res 161:212–219
- Mesnage R, Bernay B, Séralini GE (2013) Ethoxylated adjuvants of glyphosate-based herbicides are active principles of human cell toxicity. Toxicology 313(2–3):122–128
- Mineur YS, Obayemi A, Wigestrand MB, Fote GM, Calarco CA, Li AM, Picciotto MR (2013) Cholinergic signaling in the hippocampus regulates social stress resilience and anxiety- and depressionlike behavior. Proc Natl Acad Sci 110:3573–3578
- Mitchell DG, Chapman PM, Long TJ (1987) Acute toxicity of roundup and Rodeo herbicides to rainbow trout, chinook, and coho salmon. Bull Environ Contam Toxicol 39(6):1028–1035
- Mose T, Kjaerstad MB, Mathiesen L, Nielsen JB, Edelfors S, Knudsen LE (2008) Placental passage of benzoic acid, caffeine, and glyphosate in an ex vivo human perfusion system. J Toxicol Environ Health Part A 71(15):984–991
- National Research Council (2000) Scientific frontiers in developmental toxicology and risk assessment. National Academy Press, Washington
- Nishiyori Y, Nishida M, Shioda K, Suda S, Kato S (2014) Unilateral hippocampal infarction associated with an attempted suicide: a case report. J Med Case Rep 8(1):25
- Olney JW (2002) New insights and new issues in developmental neurotoxicology. Neurotoxicology 23:659–668

- Paxinos G, Franklin KBJ (2001) The mouse brain in stereotaxic coordinates, 2nd edn. Academic Press, San Diego, p 216
- Pellow S, Chopin P, File SE, Briley M (1985) Validation of open: closed arm entries in an elevated plus-maze as a measure of anxiety in the rat. J Neurosci Methods 14(3):149–167
- Perez-Reyes E, Cribbs LL, Daud A, Lacerda AE, Barclay J, Williamson MP, Lee JH (1998) Molecular characterization of a neuronal low-voltage-activated T-type calcium channel. Nature 391(6670):896–900
- Powles SB, Preston C, Bryan IB, Jutsum AR (1997) Herbicide resistance: impact and management. Adv Agron 58:57–93
- Richard S, Moslemi S, Sipahutar H, Benachour N, Seralini GE (2005) Differential effects of glyphosate and roundup on human placental cells and aromatase. Environ Health Perspect 113(6):716–720
- Rio DC, Ares M, Hannon GJ, Nilsen TW (2010) Purification of RNA using TRIzol (TRI reagent). Cold Spring Harbor Protoc 6:5439
- Rodier PM (1995) Developing brain as a target of toxicity. Environ Health Perspect 103(6):73–76
- Rodríguez VM, Limón-Pacheco JH, Mendoza-Trejo MS, González-Gallardo A, Hernández Plata I, Giordano M (2013) Repeated exposure to the herbicide atrazine alters locomotor activity and the nigrostriatal dopaminergic system of the albino rat. Neurotoxicology 34:82–94
- Sakata K, Martinowich K, Woo NH, Schloesser RJ, Jimenez DV, Ji Y, Lu B (2013) Role of activity-dependent BDNF expression in hippocampal-prefrontal cortical regulation of behavioral perseverance. Proc Natl Acad Sci 110(37):15103–15108
- Santillán ME, Vincenti LM, Martini AC, Fiol de Cuneo M, Ruiz RD, Mangeaud A, Stutz G (2010) Developmental and neurobehavioral effects of perinatal exposure to diets with different ω-6: ω-3 ratios in mice. Nutrition 26(4):423–431
- Scheffel C, Niessen KV, Rappenglück S, Wanner KT, Thiermann H, Worek F, Seeger T (2018) Counteracting desensitization of human α7-nicotinic acetylcholine receptors with bispyridinium compounds as an approach against organophosphorus poisoning. Toxicol Lett 293:149–150
- Secher T, Novitskaia V, Berezin V, Bock E, Glenthoj B, Klementiev BA (2006) Neural cell adhesion molecule-derived fibroblast Growth factor receptor agonist, the FGL peptide, promotes early postnatal sensorymotor development and enhances social memory retention. Neurosciences 141:1289–1299
- Seneff S, Swanson N, Li C (2015) Aluminum and glyphosate can synergistically induce pineal gland pathology: connection to gut dysbiosis and neurological disease. J Agric Sci 6:42–701
- Silverman JL, Yang M, Lord C, Crawley JN (2010) Behavioural phenotyping assays for mouse models of autism. Nat Rev Neurosci 11(7):490–502
- Solomon KR (2016) Glyphosate in the general population and in applicators: a critical review of studies on exposures. Crit Rev Toxicol 46:21–27
- Solomon KR, Anadón A, Carrasquilla G, Cerdeira AL, Marshall J, Sanin LH (2007) Coca and poppy eradication in Colombia: environmental and human health assessment of aerially applied glyphosate. Rev Environ Contam Toxicol 190:43–125
- Thiruchelvam M, Richfield EK, Baggs RB, Tank AW, Cory-Slechta DA (2000) The nigrostriatal dopaminergic system as a preferential target of repeated exposures to combined paraquat and maneb: implications for Parkinson's disease. J Neurosci 20(24):9207–9214
- Voorhees JR, Rohlman DS, Lein PJ, Pieper AA (2017) Neurotoxicity in preclinical models of occupational exposure to organophosphorus compounds. Front Neurosci 10:590
- Walsh RN, Cummins RA (1976) The open-field test: a critical review. Psychol Bull 83(3):482
- Walsh LP, McCormick C, Martin C, Stocco DM (2000) Roundup inhibits steroidogenesis by disrupting steroidogenic acute

regulatory (StAR) protein expression. Environ Health Perspect 108(8):769–776

- Wang G, Fan XN, Tan YY, Cheng Q, Chen SD (2011) Parkinsonism after chronic occupational exposure to glyphosate. Parkinsonism Relat Disord 17:486–487
- World Health Organisation (WHO), United Nations Environment Programme, the International Labour Organisation. (1994). Glyphosate. Environmental Health Criteria #159. Geneva, Switzerland: World Health Organization
- Yoon T, Okada J, Jung MW, Kim JJ (2008) Prefrontal cortex and hippocampus subserve different components of working memory in rats. Learn Mem 15:97–105

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.